

Quasi-invariant and pseudo-differentiable measures on a non-Archimedean Banach space.I. Real-valued measures.

Sergey V. Ludkovsky.

09.05.2001

Abstract

Quasi-invariant and pseudo-differentiable measures on a Banach space X over a non-Archimedean locally compact infinite field with a non-trivial valuation are defined and constructed. Measures are considered with values in $\mathbf{R}$. Theorems and criteria are formulated and proved about quasi-invariance and pseudo-differentiability of measures relative to linear and non-linear operators on X . Characteristic functionals of measures are studied. Moreover, the non-Archimedean analogs of the Bochner-Kolmogorov and Minlos-Sazonov theorems are investigated. Infinite products of measures also are considered. Convergence of quasi-invariant and pseudo-differentiable measures in the corresponding spaces of measures is investigated.

1 Introduction.

There are few works about integration in a classical Banach space, that is over the field $\mathbf{R}$ of real numbers or the field $\mathbf{C}$ of complex numbers [5, 6, 7, 8, 38, 42]. On the other hand, for a non-Archimedean Banach space X (that is over a non-Archimedean field) this theory is less developed. An integration in X is a very important part of the non-Archimedean analysis. The progress of quantum mechanics and different branches of modern physics related, for example, with theories of elementary particles lead

to the necessity of developing integration theory in a non-Archimedean Banach space [1, 9, 16, 43, 17]. It may also be useful for the development of non-Archimedean analysis. Non-Archimedean functional analysis develops rapidly in recent years and has many principal differences from the classical functional analysis [18, 34, 35, 36, 37, 43]. Topological vector spaces over non-Archimedean fields are totally disconnected, classes of smoothness for functions and compact operators are defined for them quite differently from that of the classical case, also the notion of the orthogonality of vectors has obtained quite another meaning. In the non-Archimedean case analogs of the Radon-Nikodym theorem and the Lebesgue theorem about convergence are true under more rigorous and another conditions. Especially strong differences are for measures with values in non-Archimedean fields, because classical notions of σ -additivity and quasi-invariance have lost their meaning.

On the other hand the development of the non-Archimedean functional analysis and its applications in non-Archimedean quantum mechanics [43, 18, 17] leads to the necessity of solving such problems. For example, problems related with quantum mechanics on manifolds are related with diffeomorphism groups and loop groups, their representations and measures on them [16, 22, 26]. In articles [20, 22, 24, 26] quasi-invariant measures on diffeomorphism and loop groups and also on non-Archimedean manifolds were constructed. These measures were used for the investigation of irreducible representations of topological groups [22, 24, 25]. The theorems proved in this work enlarge classes of measures on such groups and manifolds, this also enlarges classes of representations. For example, theorems of the Minlos-Sazonov type characterize measures with the help of characteristic functionals and compact operators. In the non-Archimedean case compact operators are more useful, than nuclear operators in the classical case. Theorems of the Bochner-Kolmogorov and Kakutani type characterize products of measures and their absolute continuity relative to others measures.

In this article measures are considered on Banach spaces, though the results given below can be developed for more general topological vector spaces, for example, it is possible to follow the ideas of works [29, 30, 31], in which were considered non-Archimedean analogs of the Minlos-Sazonov theorems for real-valued measures on topological vector spaces over non-Archimedean fields of zero characteristic. But it is impossible to make in one article. In this article, apart from articles of Mądrecki, measures are considered also with

values in non-Archimedean fields, for the cases of real-valued measures also Banach spaces over non-Archimedean fields $\mathbf{K}$ of characteristic $\text{char}(\mathbf{K}) > 0$ are considered. It is well-known, that a real-valued measure m on a locally compact Hausdorff totally disconnected Abelian topological group G is called the Haar measure, if

$$(H) \quad m(x + A) = m(A) \text{ for each } x \in G \text{ and each Borel subset } A \text{ in } G.$$

For the s -free group G a measure m with values in a non-Archimedean field $\mathbf{K}_s$ satisfy condition (H) only for an algebra of clopen (closed and open) subsets A , where a field $\mathbf{K}_s$ is a finite algebraic extension of $\mathbf{Q}_s$. Indeed, in the last case if a measure is locally finite and σ -additive on the Borel algebra of G , then it is purely atomic with atoms being singletons, so it can not be invariant relative to the entire Borel algebra (see Chapters 7-9 [34]). A lot of definitions and theorems given below are the non-Archimedean analogs of classical results. Frequently their formulations and proofs differ strongly. If proofs differ slightly, only general circumstances are given in the non-Archimedean case.

In §2 weak distributions, characteristic functions of measures and their properties are defined and investigated. The non-Archimedean analogs of the Minlos-Sazonov and Bochner-Kolmogorov theorems are given. Quasi-measures also are considered. In §3 products of measures are considered together with their density functions. In the present paper broad classes of quasi-invariant measures are defined and constructed. Theorems about quasi-invariance of measures under definite linear and non-linear transformations $U : X \rightarrow X$ are proved. §4 contains a notion of pseudo-differentiability of measures. This is necessary, because for functions $f : \mathbf{K} \rightarrow \mathbf{R}$ there is not any notion of differentiability (there is not such non-linear non-trivial f). There are given criteria for the pseudo-differentiability. In §5 there are given theorems about convergence of measures with taking into account their quasi-invariance and pseudo-differentiability, that is, in the corresponding spaces of measures. The main results are Theorems 2.27, 2.35, 3.4, 3.20, 3.24, 3.25, 4.2, 4.3, 4.5, 4.7, 5.7-5.10.

In the first part of this article real-valued measures are considered. In the second part measures with values in non-Archimedean fields are considered. This is caused by differences in definitions, formulations of statements and their proofs in two such cases. When differences are small only briefly matters are discussed in the second case.

Notations. Henceforth, $\mathbf{K}$ denotes a locally compact infinite field with

a non-trivial norm, then the Banach space X is over $\mathbf{K}$. In the present article measures on X have values in $\mathbf{R}$. We assume that either $\mathbf{K}$ is a finite algebraic extension of the field of p -adic numbers $\mathbf{Q}_p$ or $\text{char}(\mathbf{K}) = p$ and $\mathbf{K}$ is isomorphic with a field $\mathbf{F}_p(\theta)$ of formal power series consisting of elements $x = \sum_j a_j \theta^j$, where $a_j \in \mathbf{F}_p$, $|\theta| = p^{-1}$, $\mathbf{F}_p$ is a finite field of p elements, p is a prime number. These imply that $\mathbf{K}$ has the Haar measures with values in $\mathbf{R}$ [34]. If X is a Hausdorff topological space with a small inductive dimension $\text{ind}(X) = 0$, then

E denotes an algebra of subsets of X , as a rule $E \supset Bf(X)$ for real-valued measures, where

$Bf(X)$ is a Borel σ -field of X in §2.1;

$Af(X, \mu)$ is the completion of E by a measure μ in §2.1;

$M(X)$ is a space of norm-bounded measures on X in §2.1;

$M_t(X)$ is a space of Radon norm-bounded measures in §2.1;

$c_0(\alpha, \mathbf{K})$ is a Banach space and P_L is a projector (fixed relative to a chosen basis) in §2.2;

μ_L is a projection of a measure μ in §2.2;

$\{\mu_{L_n} : n\}$ is a sequence of weak distributions in §2.2;

$B(X, x, r)$ is a ball in §2.2;

χ_ξ is a character with values in $\mathbf{T} \subset \mathbf{C}$ in §2.6;

$\theta(z) = \hat{\mu}$ is a characteristic functional in §2.6;

δ_0 is the Dirac measure in §2.8;

$\mu_1 * \mu_2$ is a convolution of measures in §2.11;

$\psi_{q, \mu}$ and τ_q in §2.14;

$C(X, \mathbf{K})$ is a space of continuous functions from X into $\mathbf{K}$ in §2.6;

X^* is the topological dual space of X [33];

$\hat{C}(Y, \Gamma)$, $\tau(Y)$ in §2.26;

B_+ , C_+ in §2.29;

$\nu \ll \mu$, $\nu \sim \mu$, $\nu \perp \mu$ in §2.36.

2 Weak distributions and families of measures.

2.1. For a Hausdorff topological space X with a small inductive dimension $\text{ind}(X) = 0$ [10] the Borel σ -field is denoted $Bf(X)$. Henceforth, measures μ are given on a measurable space (X, E) . The completion of E relative to μ is denoted by $Af(X, \mu)$. The total variation of μ with values in $\mathbf{R}$ on a subset

A is denoted by $\|\mu|_A\|$ or $|\mu|(A)$ for $A \in Af(X, \mu)$. If μ is non-negative and $\mu(X) = 1$, then it is called a probability measure.

A measure μ on E is called Radon, if for each $\epsilon > 0$ there exists a compact subset $C \subset X$ such that $\|\mu|_{(X \setminus C)}\| < \epsilon$. Henceforth, $\mathbf{M}(X)$ denotes a space of norm-bounded measures, $\mathbf{M}_t(X)$ is its subspace of Radon norm-bounded measures.

2.2. Each Banach space X over $\mathbf{K}$ in view of Theorems 5.13 and 5.16 [34] is isomorphic with $c_0(\alpha, \mathbf{K}) := \{x : x = (x_j : j \in \alpha), \text{card}(j : |x_j|_{\mathbf{K}} > b) < \aleph_0 \text{ for each } b > 0\}$, where α is a set considered as an ordinal due to the Kuratowski-Zorn lemma, $\text{card}(A)$ denotes the cardinality of A , $\|x\| := \sup(|x_j| : j \in \alpha)$. A dimension of X over $\mathbf{K}$ is by the definition $\dim_{\mathbf{K}} X := \text{card}(\alpha)$. For each closed $\mathbf{K}$ -linear subspace L in X there exists an operator of a projection $P_L : X \rightarrow L$. Moreover, an orthonormal in the non-Archimedean sense basis in L has a completion to an orthonormal basis in X such that P_L can be defined in accordance with a chosen basis.

If $A \in Bf(L)$, then $P_L^{-1}(A)$ is called a cylindrical subset in X with a base A , $B^L := P_L^{-1}(Bf(L))$, $B_0 := \cup(B^L : L \subset X, L \text{ is a Banach subspace}, \dim_{\mathbf{K}} X < \aleph_0)$. The minimal σ -algebra σB_0 generated by B_0 coincides with $Bf(X)$, if $\dim_{\mathbf{K}} X \leq \aleph_0$. Henceforward, it is assumed that $\alpha \leq \omega_0$, where ω_0 is the initial ordinal with the cardinality $\aleph_0 := \text{card}(\mathbf{N})$. Then there exists an increasing sequence of Banach subspaces $L_n \subset L_{n+1} \subset \dots$ such that $cl(\cup[L_n : n]) = X$, $\dim_{\mathbf{K}} L_n = \kappa_n$ for each n , where $cl(A) = \bar{A}$ denotes a closure of A in X for $A \subset X$. We fix a family of projections $P_{L_n}^{L_m} : L_m \rightarrow L_n$ such that $P_{L_n}^{L_m} P_{L_k}^{L_n} = P_{L_k}^{L_m}$ for each $m \geq n \geq k$. A projection of the measure μ onto L denoted by $\mu_L(A) := \mu(P_L^{-1}(A))$ for each $A \in Bf(L)$ compose the consistent family:

$$(1) \mu_{L_n}(A) = \mu_{L_m}(P_{L_n}^{-1}(A) \cap L_m)$$

for each $m \geq n$, since there are projectors $P_{L_n}^{L_m}$, where $\kappa_n \leq \aleph_0$ and there may be chosen $\kappa_n < \aleph_0$ for each n .

An arbitrary family of measures $\{\mu_{L_n} : n \in \mathbf{N}\}$ having property (1) is called a sequence of a weak distribution (see also [8, 38]).

By $B(X, x, r)$ we denote a ball $\{y : y \in X, \|x - y\| \leq r\}$, which is clopen (closed and open) in X .

2.3. Lemma. *A sequence of a weak distribution $\{\mu_{L_n} : n\}$ is generated by some measure μ on $Bf(X)$ if and only if for each $c > 0$ there exists $b > 0$ such that $|\mu_{L_n}|(B(X, 0, r) \cap L_n) - |\mu_{L_n}|(L_n)| \leq c$ and $\sup_n |\mu_{L_n}|(L_n) < \infty$.*

Proof. In the case of μ with values in $\mathbf{R}$ we can use a Hahn decomposition $\mu = \mu^+ - \mu^-$ and substitute everywhere in the proof of Lemma 1 §2[38] a Hilbert space over $\mathbf{R}$ onto the Banach space X over $\mathbf{K}$, since X is the Radon space in view of Theorem 1.2 §I.1.3 [8], then $|\mu|(A) = \mu^+(A) + \mu^-(A)$ for $A \in Bf(X)$.

2.4. Definition and notations. A function $\phi : X \rightarrow \mathbf{R}$ of the form $\phi(x) = \phi_S(P_S x)$ is called a cylindrical function if ϕ_S is a $Bf(S)$ -measurable function on a finite-dimensional over $\mathbf{K}$ space S in X . For $\phi_S \in L^1(S, \mu, \mathbf{R})$ for μ with values in $\mathbf{R}$ we can define an integral by a sequence of a weak distribution $\{\mu_{S_n} : n\}$:

$$\int_X \phi(x) \mu_*(dx) := \int \phi_{S_n}(x) \mu_{S_n}(dx).$$

2.5. Lemma. A subset $A \subset X = c_0(\omega_0, \mathbf{K})$ is relatively compact if and only if A is bounded and for each $c > 0$ there exists a finite-dimensional over $\mathbf{K}$ subspace $L \subset X$ such that $\bar{A} \subset L^c := \{y \in X : d(y, L) := \inf\{\|x - y\| : x \in L\} \leq c\}$.

Proof. If A is bounded and for each $c > 0$ there exists L^c with $\bar{A} \subset L^c$, then there is a sequence $\{k(j) : j \in \mathbf{N}\} \subset \mathbf{Z}$ such that $\lim_{j \rightarrow \infty} k(j) = \infty$, $\bar{A} \subset \{x \in X : |x_j| \leq p^{-k(j)}, j = 1, 2, \dots\} =: S$, but X is Lindelöf, S is sequentially compact, hence $\bar{A}$ is compact (see §3.10.31 [10]). If $\bar{A}$ is compact, then for each $c > 0$ there exists a finite number m such that $\bar{A} \subset \bigcup_{j=1}^m B(X, x_j, c)$, where $x_j \in X$. Therefore, $\bar{A} \subset L^c$ for $L = sp_{\mathbf{K}}(x_j : j = 1, \dots, m) := (x = \sum_{j=1}^m b_j x_j : b_j \in K)$.

2.6. Remarks and definitions. When $char(\mathbf{K}) = 0$, then $\mathbf{K}$ as the $\mathbf{Q}_p$ linear space is isomorphic with $\mathbf{Q}_p^n$, where $n \in \mathbf{N} := \{1, 2, \dots\}$. The topologically adjoint space over $\mathbf{Q}_p$ (that is, of continuous linear functionals $f : \mathbf{K} \rightarrow \mathbf{Q}_p$) is isomorphic with $\mathbf{Q}_p^n$ [15]. For x and $z \in \mathbf{Q}_p^n$ we denote by (z, x) the following sum $\sum_{j=1}^n x_j z_j$, where $x = (x_j : j = 1, \dots, n)$, $x_j \in \mathbf{Q}_p$. Each number $y \in \mathbf{Q}_p$ has a decomposition $y = \sum_l a_l p^l$, where $\min(l : a_l \neq 0) =: ord_p(y) > -\infty$ ($ord(0) := \infty$) [33], $a_l \in (0, 1, \dots, p-1)$, we define a symbol $\{y\}_p := \sum_{l < 0} a_l p^l$ for $|y|_p > 1$ and $\{y\}_p = 0$ for $|y|_p \leq 1$.

For a locally compact field $\mathbf{K}$ with a characteristic $char(\mathbf{K}) = p > 0$ let $\pi_j(x) := a_j$ for each $x = \sum_j a_j \theta^j \in \mathbf{K}$ (see Notation). For $\xi \in \mathbf{K}^*$ we denote $\xi(x)$ also by (ξ, x) . All continuous characters $\chi : \mathbf{K} \rightarrow \mathbf{C}$ have the form

$$(1) \chi_\xi(x) = \epsilon^{z^{-1}\eta((\xi, x))}$$

for each $\eta((\xi, x)) \neq 0$, $\chi_\xi(x) := 1$ for $\eta((\xi, x)) = 0$, where $\epsilon = 1^z$ is a root of unity, $z = p^{\text{ord}(\eta((\xi, x)))}$, $\pi_j : \mathbf{K} \rightarrow \mathbf{R}$, $\eta(x) := \{x\}_p$ and $\xi \in \mathbf{Q}_p^{n*} = \mathbf{Q}_p^n$ for $\text{char}(\mathbf{K}) = 0$, $\eta(x) := \pi_{-1}(x)/p$ and $\xi \in \mathbf{K}^* = \mathbf{K}$ for $\text{char}(\mathbf{K}) = p > 0$, $x \in \mathbf{K}$, (see §25 [15]). Each χ is locally constant, hence $\chi : \mathbf{K} \rightarrow \mathbf{T}$ is also continuous, where $\mathbf{T}$ denotes the discrete group of all roots of 1 (by multiplication).

For a measure μ there exists a characteristic functional (that is, called the Fourier-Stieltjes transformation) $\theta = \theta_\mu : C(X, \mathbf{K}) \rightarrow \mathbf{C}$:

$$(2) \theta(f) := \int_X \chi_e(f(x)) \mu(dx),$$

where either $e = (1, \dots, 1) \in \mathbf{Q}_p^n$ for $\text{char}(\mathbf{K}) = 0$, or $e = 1 \in \mathbf{K}^*$ for $\text{char}(\mathbf{K}) = p > 0$, $x \in X$, f is in the space $C(X, \mathbf{K})$ of continuous functions from X into $\mathbf{K}$, in particular for $z = f$ in the topologically conjugated space X^* over $\mathbf{K}$, $z : X \rightarrow \mathbf{K}$, $z \in X^*$, $\theta(z) =: \hat{\mu}(z)$. It has the following properties:

$$(3a) \theta(0) = 1 \text{ for } \mu(X) = 1$$

and $\theta(f)$ is bounded on $C(X, \mathbf{K})$;

$$(3b) \sup_f |\theta(f)| = 1 \text{ for probability measures ;}$$

(4) $\theta(z)$ is weakly continuous, that is, $(X^*, \sigma(X^*, X))$ -continuous,

$\sigma(X^*, X)$ denotes a weak topology on X^* , induced by the Banach space X over $\mathbf{K}$. To each $x \in X$ there corresponds a continuous linear functional $x^* : X^* \rightarrow \mathbf{K}$, $x^*(z) := z(x)$, moreover, $\theta(f)$ is uniformly continuous relative to the norm on

$$C_b(X, \mathbf{K}) := \{f \in C(X, \mathbf{K}) : \|f\| := \sup_{x \in X} |f(x)|_{\mathbf{K}} < \infty\};$$

(5) $\theta(z)$ is positive definite on X^* and on $C(X, \mathbf{K})$

for μ with values in $[0, \infty)$.

Property (4) follows from Lemma 2.3, boundedness and continuity of χ_e and the fact that due to the Hahn-Banach theorem there is $x_z \in X$ with $z(x_z) = 1$ for $z \neq 0$ such that $z|_{(X \ominus L)} = 0$ and

$$\theta(z) = \int_X \chi_e(P_L(x)) \mu(dx) = \int_L \chi_e(y) \mu_L(dy),$$

where $L = Kx_z$, also due to the Lebesgue theorem 2.4.9 [11] for real measures. Indeed, for each $c > 0$ there exists a compact subset $S \subset X$ such that $|\mu|(X \setminus S) < c$, each bounded subset $A \subset X^*$ is uniformly equicontinuous on S (see (9.5.4) and Exer. 9.202 [33]), that is, $\{\chi_e(z(x)) : z \in A\}$ is the uniformly equicontinuous family (by $x \in S$). On the other hand, $\chi_e(f(x))$ is uniformly equicontinuous on a bounded $A \subset C_b(X, \mathbf{K})$ by $x \in S$.

Property (5) is accomplished, since

$$\sum_{l,j=1}^N \theta(f_l - f_j) \alpha_l \bar{\alpha}_j = \int_X \left| \sum_{j=1}^N \alpha_j \chi_e(f_j(x)) \right|^2 \mu(dx) \geq 0,$$

particularly, for $f_j = z_j \in X$, where $\bar{\alpha}_j$ is a complex conjugated number to α_j .

We call a functional θ finite-dimensionally concentrated, if there exists $L \subset X$, $\dim_{\mathbf{K}} L < \aleph_0$, such that $\theta|_{(X \setminus L)} = \mu(X)$. For each $c > 0$ and $\delta > 0$ in view of Theorem I.1.2 [8] and Lemma 2.5 there exists a finite-dimensional over $\mathbf{K}$ subspace L and compact $S \subset L^\delta$ such that $\|X \setminus S\|_\mu < c$. Let $\theta^L(z) := \theta(P_L z)$.

This definition is correct, since $L \subset X$, X has the isometrical embedding into X^* as the normed space associated with the fixed basis of X , such that functionals $z \in X$ separate points in X . If $z \in L$, then $|\theta(z) - \theta^L(z)| \leq c \times b \times q$, where $b = \|X\|_\mu$, q is independent of c and b . Each characteristic functional $\theta^L(z)$ is uniformly continuous by $z \in L$ relative to the norm $\|\cdot\|$ on L , since $|\theta^L(z) - \theta^L(y)| \leq |\int_{S' \cap L} [\chi_e(z(x)) - \chi_e(y(x))] \mu_L(dx)| + |\int_{L \setminus S'} [\chi_e(z(x)) - \chi_e(y(x))] \mu_L(dx)|$, where the second term does not exceed $2C'$ for $\|L \setminus S'\|_{\mu_L} < c'$ for a suitable compact subset $S' \subset X$ and $\chi_e(z(x))$ is an uniformly equicontinuous by $x \in S'$ family relative to $z \in B(L, 0, 1)$.

Therefore,

$$(6) \quad \theta(z) = \lim_{n \rightarrow \infty} \theta_n(z)$$

for each finite-dimensional over $\mathbf{K}$ subspace L , where $\theta_n(z)$ is uniformly equicontinuous and finite-dimensionally concentrated on $L_n \subset X$, $z \in X$, $cl(\bigcup_n L_n) = X$, $L_n \subset L_{n+1}$ for every n , for each $c > 0$ there are n and $q > 0$ such that $|\theta(z) - \theta_j(z)| \leq cbq$ for $z \in L_j$ and $j > n$, $q = \text{const} > 0$ is independent of j , c and b . Let $\{e_j : j \in \mathbf{N}\}$ be the standard orthonormal basis in X , $e_j = (0, \dots, 0, 1, 0, \dots)$ with 1 in j -th place. Using countable additivity of μ , local constantness of χ_e , considering all $z = be_j$ and $b \in \mathbf{K}$, we get

that $\theta(z)$ on X is non-trivial, whilst μ is a non-zero measure, since due to Lemma 2.3 μ is characterized uniquely by $\{\mu_{L_n}\}$. Indeed, for μ with values in $\mathbf{R}$ a measure μ_V on V , $\dim_{\mathbf{K}} V < \aleph_0$, this follows from the properties of the Fourier transformation F on spaces of generalized functions and also on $L^2(V, \mu_V, \mathbf{C})$ (see §7 [43]), where

$$F(g)(z) := \lim_{r \rightarrow \infty} \int_{B(V, 0, r)} \chi_e(z(x)) g(x) m(dx),$$

$z \in V$, m is the Haar measure on V with values in $\mathbf{R}$. Therefore, the mapping $\mu \mapsto \theta_\mu$ is injective.

2.7. Proposition. *Let $X = \mathbf{K}^j$, $j \in \mathbf{N}$,*

(a) μ and ν be real probability measures on X , suppose ν is symmetric. Then $\int_X \hat{\mu}(x) \nu(dx) = \int_X \hat{\nu}(x) \mu(dx) \in \mathbf{R}$ and for each $0 < l < 1$ is accomplished the following inequality:

$$\mu([x \in X : \hat{\nu}(x) \leq l]) \leq \int_X (1 - \hat{\mu}(x)) \nu(dx) / (1 - l).$$

(b). For each real probability measure μ on X there exists $r > p^3$ such that for each $R > r$ and $t > 0$ the following inequality is accomplished:

$$\mu([x \in X : \|x\| \geq tR]) \leq c \int_X [1 - \hat{\mu}(y\xi)] \nu(dy),$$

where $\nu(dx) = C \times \exp(-|x|^2) m(dx)$, m is the Haar measure on X with values in $[0, \infty)$, $m(B(X, 0, 1)) = 1$, $\nu(X) = 1$, $2 > c = \text{const} \geq 1$ is independent from t , $c = c(r)$ is non-increasing whilst r is increasing, $C > 0$.

Proof. (a). Recall that ν is symmetric, if $\nu(B) = \nu(-B)$ for each $B \in Bf(X)$. Therefore, $\int_X \chi_e(z(x)) \nu(dx) = \int_X \chi_e(-z(x)) \nu(dx)$, that is equivalent to $\int_X [\chi_e(z(x)) - \chi_e(-z(x))] \nu(dx) = 0$ or $\hat{\nu}(z) \in \mathbf{R}$. If $0 < l < 1$, then $\mu([x \in X : \hat{\nu}(x) \leq l]) = \mu([x : 1 - \hat{\nu}(x) \geq 1 - l]) \leq \int_X (1 - \hat{\nu}(x)) \mu(dx) / (1 - l) = \int_X (1 - \hat{\mu}(x)) \nu(dx) / (1 - l)$ due to the Fubini theorem.

(b). Let $\nu(dx) = \gamma(x) m(dx)$, where $\gamma(x) = C \times \exp(-|x|^2)$, $C > 0$, $\nu(X) = 1$. Then $F(\gamma)(z) =: \hat{\gamma}(z) \geq 0$, and $\hat{\gamma}(0) = 1$ and γ is the continuous positive definite function with $\gamma(z) \rightarrow 0$ whilst $|z| \rightarrow \infty$. In view of (a): $\mu([x : \|x\| \geq tR]) \leq \int_X [1 - \hat{\mu}(y\xi)] \nu(dy) / (1 - l)$, where $|\xi| = 1/t$, $t > 0$, $l = l(R)$. Estimating integrals, we get (b).

2.8. Lemma. *Let in the notation of Proposition 2.7 $\nu_\xi(dx) = \gamma_\xi(x) m(dx)$, $\gamma_\xi(x) = C(\xi) \exp(-|x\xi|^2)$, $\nu_\xi(X) = 1$, $\xi \neq 0$, then a measure ν_ξ is weakly converging to the Dirac measure δ_0 with the support in $0 \in X$ for $|\xi| \rightarrow \infty$.*

Proof. We have: $C(\xi)^{-1} = C_q(\xi)^{-1} = \sum_{l \in \mathbf{Z}} [p^{lq} - p^{(l-1)q}] \exp(-p^{2l} |\xi|^2) < \infty$, where the sum by $l < 0$ does not exceed 1, $q = jn$, $j = \dim_{\mathbf{K}} X$, $n = \dim_{\mathbf{Q}_p} \mathbf{K}$. Here $\mathbf{K}$ is considered as the Banach space $\mathbf{Q}_p^n$ with the

following norm $|\cdot|_p$ equivalent to $|\cdot|_{\mathbf{K}}$, for $x = (x_1, \dots, x_j) \in X$ with $x_l \in \mathbf{K}$ as usually $|x|_p = \max_{1 \leq l \leq j} |x_l|_p$, for $y = (y_1, \dots, y_n) \in \mathbf{K}$ with $y_l \in \mathbf{Q}_p$: $|y|_p := \max_{1 \leq l \leq n} |y_l|_{\mathbf{Q}_p}$. Further, $p^{l+s} \sum_{x_l \neq 0} \exp(2\pi i \sum_{i=l}^{-s-1} x_i p^{i+s}) = \int_{p^{l+s}}^1 \exp(2\pi i \phi) d\phi + \beta(s)$, where $s + l < 0$, $\lim_{s \rightarrow -\infty} (\beta(s) p^{-s-l}) = 0$, therefore, $\sup[|\hat{\gamma}_1(z)|_{\mathbf{R}} |z|_X : z \in X, |z| \geq p^3] \leq 2$. Then taking $0 \neq \xi \in \mathbf{K}$ and carrying out the substitution of variable for continuous and bounded functions $f : X \rightarrow \mathbf{R}$ we get $\lim_{|\xi| \rightarrow \infty} \int_X f(x) \nu_\xi(dx) = f(0)$. This means that ν_ξ is weakly converging to δ_0 for $|\xi| \rightarrow \infty$.

2.9. Theorem. *Let μ_1 and μ_2 be measures in $\mathbf{M}(X)$ such that $\hat{\mu}_1(f) = \hat{\mu}_2(f)$ for each $f \in \Gamma$. Then $\mu_1 = \mu_2$, where $X = c_0(\alpha, K)$, $\alpha \leq \omega_0$, Γ is a vector subspace in a space of continuous functions $f : X \rightarrow \mathbf{K}$ separating points in X .*

Proof. Let at first $\alpha < \omega_0$, then due to continuity of the convolution $\gamma_\xi * \mu_j$ by ξ , and Proposition 4.5 §I.4[42] and Lemma 2.8 we get $\mu_1 = \mu_2$, since the family Γ generates $Bf(X)$. Now let $\alpha = \omega_0$, $A = \{x \in X : (f_1(x), \dots, f_n(x)) \in S\}$, ν_j be an image of a measure μ_j for a mapping $x \mapsto (f_1(x), \dots, f_n(x))$, where $S \in Bf(\mathbf{K}^n)$, $f_j \in X \hookrightarrow X^*$. Then $\hat{\nu}_1(y) = \hat{\mu}_1(y_1 f_1 + \dots + y_n f_n) = \hat{\mu}_2(y_1 f_1 + \dots + y_n f_n) = \hat{\nu}_2(y)$ for each $y = (y_1, \dots, y_n) \in \mathbf{K}^n$, consequently, $\nu_1 = \nu_2$ on E . Further we can use the Prohorov theorem 3.4 §1.3 [42], since compositions of $f \in \Gamma$ with continuous functions $g : \mathbf{K} \rightarrow \mathbf{R}$ generate a family of real-valued functions separating points of X .

2.10. Proposition. *Let μ_l and μ be measures in $\mathbf{M}(X_l)$ and $\mathbf{M}(X)$ respectively, where $X_l = c_0(\alpha_l, \mathbf{K})$, $\alpha_l \leq \omega_0$, $X = \prod_{l=1}^n X_l$, $n \in \mathbf{N}$. Then the condition $\hat{\mu}(z_1, \dots, z_n) = \prod_{l=1}^n \hat{\mu}_l(z_l)$ for each $(z_1, \dots, z_n) \in X \hookrightarrow X^*$ is equivalent to $\mu = \prod_{l=1}^n \mu_l$.*

Proof. Let $\mu = \prod_{l=1}^n \mu_l$, then $\hat{\mu}(z_1, \dots, z_n) = \int_X \chi_e(\sum z_l(x_l)) \prod_{l=1}^n \mu_l(dx_l) = \prod_{l=1}^n \int_{X_l} \chi_e(z_l(x_l)) \mu_l(dx_l)$. The reverse statement follows from Theorem 2.9.

2.11. Proposition. *Let X be a Banach space over $\mathbf{K}$; suppose μ , μ_1 and μ_2 are probability measures on X . Then the following conditions are equivalent: μ is the convolution of two measures μ_j , $\mu = \mu_1 * \mu_2$, and $\hat{\mu}(z) = \hat{\mu}_1(z) \hat{\mu}_2(z)$ for each $z \in X$.*

Proof. Let $\mu = \mu_1 * \mu_2$. This means by the definition that μ is the image of the measure $\mu_1 \otimes \mu_2$ for the mapping $(x_1, x_2) \mapsto x_1 + x_2$, $x_j \in X$, consequently, $\hat{\mu}(z) = \int_{X \times X} \chi_e(z(x_1 + x_2)) (\mu_1 \otimes \mu_2)(d(x_1, x_2)) = \prod_{l=1}^2 \int_X \chi_e(z(x_l)) \mu_l(dx_l) = \hat{\mu}_1(z) \hat{\mu}_2(z)$. On the other hand, if $\hat{\mu}_1 \hat{\mu}_2 = \hat{\mu}$, then $\hat{\mu} = (\mu_1 * \mu_2)^\wedge$ and due to Theorem 2.9 above for real measures we have $\mu = \mu_1 * \mu_2$.

2.12. Corollary. *Let ν be a probability measure on $Bf(X)$ and $\mu * \nu = \mu$ for each μ , then $\nu = \delta_0$.*

Proof. If $z_0 \in X \hookrightarrow X^*$ and $\hat{\mu}(z_0) \neq 0$, then from $\hat{\mu}(z_0)\hat{\nu}(z_0) = \hat{\mu}(z_0)$ it follows that $\hat{\nu}(z_0) = 1$. From the property 2.6(6) we get that there exists $m \in \mathbf{N}$ with $\hat{\mu}(z) \neq 0$ for each z with $\|z\| = p^{-m}$, since $\hat{\mu}(0) = 1$. Then $\hat{\nu}(z + z_0) = 1$, that is, $\hat{\nu}|_{(B(X, z_0, p^{-m}))} = 1$. Since μ are arbitrary we get $\hat{\nu}|_X = 1$, that is, $\nu = \delta_0$ due to §2.6 and §2.9.

2.13. Corollary. *Let X and Y be Banach spaces over $\mathbf{K}$, (a) μ and ν be probability measures on X and Y respectively, suppose $T : X \rightarrow Y$ is a continuous linear operator. A measure ν is an image of μ for T if and only if $\hat{\nu} = \hat{\mu} \circ T^*$, where $T^* : Y^* \rightarrow X^*$ is an adjoint operator. (b). A characteristic functional of a real measure μ on $Bf(X)$ is real if and only if μ is symmetric.*

Proof follows from §2.6 and §2.9.

2.14. Definition. We say that a real probability measure μ on $Bf(X)$ for a Banach space X over $\mathbf{K}$ and $0 < q < \infty$ has a weak q -th order if $\psi_{q,\mu}(z) = \int_X |z(x)|^q \mu(dx) < \infty$ for each $z \in X^*$. The weakest vector topology in X^* relative to which all $(\psi_{q,\mu} : \mu)$ are continuous is denoted by τ_q .

2.15. Theorem. *A characteristic functional $\hat{\mu}$ of a real probability Radon measure μ on $Bf(X)$ is continuous in the topology τ_q for each $q > 0$.*

Proof. For each $c > 0$ there exists a compact $S \subset X$ such that $\mu(S) > 1 - c/4$ and

$$|1 - \hat{\mu}(z)| \leq \left| \int_S (1 - \chi_e(z(x))) \mu(dx) \right| + \left| \int_{X \setminus S} (1 - \chi_e(z(x))) \mu(dx) \right| \leq |1 - \hat{\mu}_c(z)| + c/2,$$

where $\mu_c(A) = (\mu(A \cap S))/\mu(S)$ and $A \in Bf(X)$; further analogously to the proof of IV.2.3[42].

2.16. Proposition. *For a completely regular space X with $\text{ind}(X) = 0$ the following statements are accomplished:*

(a) *if (μ_β) is a bounded net of measures in $\mathbf{M}(X)$ that weakly converges to a measure μ in $\mathbf{M}(X)$, then $(\hat{\mu}_\beta(f))$ converges to $\hat{\mu}(f)$ for each continuous $f : X \rightarrow \mathbf{K}$; if X is separable and metrizable then $(\hat{\mu}_\beta)$ converges to $\hat{\mu}$ uniformly on subsets that are uniformly equicontinuous in $C(X, \mathbf{K})$;*

(b) *if M is a bounded dense family in a ball of the space $\mathbf{M}(X)$ for measures in $\mathbf{M}(X)$, then a family $(\hat{\mu} : \mu \in M)$ is equicontinuous on a locally $\mathbf{K}$ -convex space $C(X, \mathbf{K})$ in a topology of uniform convergence on compact subsets $S \subset X$.*

Proof. (a). Functions $\chi_e(f(x))$ are continuous and bounded on X , where $\hat{\mu}(f) = \int_X \chi_e(f(x))\mu(dx)$. Then (a) follows from the definition of the weak convergence and Proposition 1.3.9[42], since $sp_{\mathbf{C}}\{\chi_e(f(x)) : f \in C(X, \mathbf{K})\}$ is dense in $C(X, \mathbf{C})$.

(b). For each $c > 0$ there exists a compact subset $S \subset X$ such that $|\mu|(S) > |\mu(X)| - c/4$. Therefore, for $\mu \in \mathbf{M}$ and $f \in C(X, \mathbf{K})$ with $|f(x)|_{\mathbf{K}} < c < 1$ for $x \in S$ we get

$$|\mu(X) - Re(\hat{\mu}(f))| = 2 \left| \int_X [(\chi_e^{1/2}(f(x)) - \chi_e^{1/2}(-f(x)))/(2i)]^2 \mu(dx) \right| < c/2.$$

Further analogously to the proof of Proposition IV.3.1[42], since X is the T_1 -space and for each point x and each closed subset S in X with $x \notin S$ there is a continuous function $h : X \rightarrow B(\mathbf{K}, 0, 1)$ such that $h(x) = 0$ and $h(S) = \{1\}$.

2.17. Theorem. *Let X be a Banach space over $\mathbf{K}$, $\eta : \Gamma \rightarrow \mathbf{C}$ be a continuous positive definite function, (μ_β) be a bounded weakly relatively compact net in the space $\mathbf{M}_t(X)$ of Radon norm-bounded measures and there exists $\lim_\beta \hat{\mu}_\beta(f) = \gamma(f)$ for each $f \in \Gamma$ and uniformly on compact subsets of the completion $\tilde{\Gamma}$, where $\Gamma \subset C(X, \mathbf{K})$ is a vector subspace separating points in X . Then (μ_β) weakly converges to $\mu \in \mathbf{M}_t(X)$ with $\hat{\mu}|_\Gamma = \gamma$.*

Proof is analogous to the proof of Theorem IV.3.1[42] and follows from Theorem 2.9 above.

2.18. Theorem. (a). *A bounded family of measures in $\mathbf{M}(\mathbf{K}^n)$ is weakly relatively compact if and only if a family $(\hat{\mu} : \mu \in M)$ is equicontinuous on $\mathbf{K}^n$.*

(b). *If $(\mu_j : j \in \mathbf{N})$ is a bounded sequence of measures in $\mathbf{M}_t(\mathbf{K}^n)$, $\gamma : \mathbf{K}^n \rightarrow \mathbf{C}$ is a continuous and positive definite function, $\hat{\mu}_j(y) \rightarrow \gamma(y)$ for each $y \in \mathbf{K}^n$, then (μ_j) weakly converges to a measure μ with $\hat{\mu} = \gamma$.*

(c). *A bounded sequence of measures (μ_j) in $\mathbf{M}_t(\mathbf{K}^n)$ weakly converges to a measure μ in $\mathbf{M}_t(\mathbf{K}^n)$ if and only if for each $y \in \mathbf{K}^n$ there exists $\lim_{j \rightarrow \infty} \hat{\mu}_j(y) = \hat{\mu}(y)$.*

(d). *If a bounded net (μ_β) in $\mathbf{M}_t(\mathbf{K}^n)$ converges uniformly on each bounded subset in $\mathbf{K}^n$, then (μ_β) converges weakly to a measure μ in $\mathbf{M}_t(\mathbf{K}^n)$, where $n \in \mathbf{N}$.*

Proof. (a). This follows from the Prohorov theorem 1.3.6[42] and Propositions 2.7, 2.16.

(b). We have the following inequality: $\lim_m \sup_{j>m} \mu_j([x \in \mathbf{K}^n : |x| \geq tR]) \leq 2 \int_{\mathbf{K}^n} (1 - \operatorname{Re}(\eta(\xi y))) \nu(dy)$ with $|\xi| = 1/t$ due to §2.7 and §2.8. In view of Theorem 2.17 (μ_j) converges weakly to μ with $\hat{\mu} = \gamma$. (c,d). These can be proved analogously to IV.3.2[42].

2.19. Corollary. *If $(\hat{\mu}_\beta) \rightarrow 1$ uniformly on some neighbourhood of 0 in $\mathbf{K}^n$ for a bounded net of measures μ_β in $\mathbf{M}_t(\mathbf{K}^n)$, then (μ_β) converges weakly to δ_0 .*

2.20. Definition. A family of probability measures $M \subset \mathbf{M}_t(X)$ for a Banach space X over $\mathbf{K}$ is called planely concentrated if for each $c > 0$ there exists a $\mathbf{K}$ -linear subspace $S \subset X$ with $\dim_{\mathbf{K}} S = n < \aleph_0$ such that $\inf(\mu(S^c) : \mu \in M) > 1 - c$. The Banach space $\mathbf{M}_t(X)$ is supplied with the following norm $\|\mu\| := |\mu|(X)$.

2.21. Lemma. *Let S and X be the same as in §2.20; $z_1, \dots, z_m \in X^*$ be a separating family of points in S . Then a set $E := S^c \cap (x \in X : |z_j(x)| \leq r_j; j = 1, \dots, m)$ is bounded for each $c > 0$ and $r_1, \dots, r_m \in (0, \infty)$.*

Proof. A space S is isomorphic with $\mathbf{K}^n$, consequently, $p(x) = \max(|z_j| : j = 1, \dots, m)$ is the norm in S equivalent to the initial norm.

2.22. Theorem. *Let X be a Banach space over $\mathbf{K}$ with a family $\Gamma \subset X^*$ separating points in $M \subset \mathbf{M}_t(X)$. Then M is weakly relatively compact if and only if a family $\{\mu_z : \mu \in M\}$ is weakly relatively compact for each $z \in \Gamma$ and M is planely concentrated, where μ_z is an image measure on $\mathbf{K}$ of a measure μ induced by z .*

Proof follows from Lemmas 2.5, 2.21 and the Prohorov theorem (see also Theorem 1.3.7[42] with a substitution $[-r_j, r_j]$ onto $B(\mathbf{K}, 0, r_j)$).

2.23. Theorem. *For X and Γ the same as in Theorem 2.22 a sequence $\{\mu_j : j \in \mathbf{N}\} \subset \mathbf{M}_t(X)$ is weakly convergent to $\mu \in \mathbf{M}_t(X)$ if and only if for each $z \in \Gamma$ there exists $\lim_{j \rightarrow \infty} \hat{\mu}_j(z) = \hat{\mu}(z)$ and a family $\{\mu_j\}$ is planely concentrated.*

Proof follows from Theorems 2.17, 18, 22 (see also Theorem IV.3.3[42]).

2.24. Proposition. *Let X be a weakly regular space with $\operatorname{ind}(X) = 0$, $\Gamma \subset C(X, \mathbf{K})$ be a vector subspace separating points in X , $(\mu_n : n \in \mathbf{N}) \subset \mathbf{M}_t(X)$, $\mu \in \mathbf{M}_t(X)$, $\lim_{n \rightarrow \infty} \hat{\mu}_n(f) = \hat{\mu}(f)$ for each $f \in \Gamma$. Then (μ_n) is weakly convergent to μ relative to the weakest topology $\sigma(X, \Gamma)$ in X relative to which all $f \in \Gamma$ are continuous.*

Proof follows from Theorem 2.18 and is analogous to the proof of Proposition IV.3.3[42].

2.25. Let $(X, \mathbf{U}) = \prod_{\lambda} (X_{\lambda}, \mathbf{U}_{\lambda})$ be a product of measurable completely

regular Radon spaces $(X_\lambda, \mathcal{U}_\lambda) = (X_\lambda, \mathcal{U}_\lambda, \mathbf{K}_\lambda)$, where $\mathbf{K}_\lambda$ are compact classes approximating from below each measure μ_λ on $(X_\lambda, \mathcal{U}_\lambda)$, that is, for each $c > 0$ and elements A of an algebra $\mathcal{U}_\lambda$ there is $S \in \mathbf{K}_\lambda$, $S \subset A$ with $\|A \setminus S\|_{\mu_\lambda} < c$.

2.26. Definition. Let X be a Banach space over $\mathbf{K}$, then a mapping $f : X \rightarrow \mathbf{C}$ is called pseudocontinuous, if its restriction $f|_L$ is uniformly continuous for each subspace $L \subset X$ with $\dim_{\mathbf{K}} L < \aleph_0$. Let Γ be a family of mappings $f : Y \rightarrow \mathbf{K}$ of a set Y into a field $\mathbf{K}$. We denote by $\hat{\mathbf{C}}(Y, \Gamma)$ the minimal σ -algebra (that is called cylindrical) generated by an algebra $\mathbf{C}(Y, \Gamma)$ of subsets of the form $C_{f_1, \dots, f_n; E} := \{x \in X : (f_1(x), \dots, f_n(x)) \in S\}$, where $S \in Bf(\mathbf{K}^n)$, $f_j \in \Gamma$. We supply Y with a topology $\tau(Y)$ which is generated by a base $(C_{f_1, \dots, f_n; E} : f_j \in \Gamma, E \text{ is open in } \mathbf{K}^n)$.

2.27. Theorem. Non-Archimedean analog of the Bochner-Kolmogorov theorem. *Let X be a Banach space over $\mathbf{K}$, X^a be its algebraically adjoint $\mathbf{K}$ -linear space (that is, of all linear mappings $f : X \rightarrow \mathbf{K}$ not necessarily continuous). A mapping $\theta : X^a \rightarrow \mathbf{C}$ is a characteristic functional of a probability measure μ with values in $\mathbf{R}$ and is defined on $\hat{\mathbf{C}}(X^a, X)$ if and only if θ satisfies conditions 2.6(3,5) for $(X^a, \tau(X^a))$ and is pseudocontinuous on X^a .*

Proof. (I). For $\dim_{\mathbf{K}} X = \text{card}(\alpha) < \aleph_0$ a space X^a is isomorphic with $\mathbf{K}^\alpha$, hence the statement of theorem for a measure μ follows from Theorems 2.9 and 2.18 above, since $\theta(0) = 1$ and $|\theta(z)| \leq 1$ for each z .

(II). We consider now the case of μ with values in $\mathbf{R}$ and $\alpha < \omega_0$. In §2.6 (see also §2.16-18,24) it was proved that $\theta = \hat{\mu}$ has the desired properties for real probability measures μ . On the other hand, there is θ which satisfies the conditions of the theorem. Let $\theta_\xi(y) = \theta(y)h_\xi(y)$, where $h_\xi(y) = F[C(\xi)\exp(-\|x\xi\|^2)](y)$ (that is, the Fourier transform by x), $\nu_\xi(\mathbf{K}^\alpha) = 1$, $\nu_\xi(dx) = C(\xi)\exp(-\|x\xi\|^2)m(dx)$ (see Lemma 2.8), $\xi \neq 0$. Then $\theta_\xi(y)$ is positive definite and is uniformly continuous as a product of two such functions. Moreover, $\theta_\xi(y) \in L^1(\mathbf{K}^\alpha, m, \mathbf{C})$. For $\xi \neq 0$ a function $f_\xi(x) = \int_{\mathbf{K}^\alpha} \theta_\xi(y)\chi_e(x(y))m(dy)$ is bounded and continuous, the function $\exp(-\|x\xi\|^2) =: s(x)$ is positive definite. Since ν_ξ is symmetric and weakly converges to δ_0 , hence there exists $r > 0$ such that for each $|\xi| > r$ we have $\hat{\gamma}_\xi(y) = \int_{\mathbf{K}^\alpha} C(\xi)\exp(-\|x\xi\|_p^2)\chi_e(y(x))m(dx) = \int [\chi_e((y(x)) + \chi_e(-y(x)))]2^{-1}\exp(-\|x\xi\|_p^2)C(\xi)m(dx)/2 > 1 - 1/R$ for $|y| \leq R$, consequently, $\hat{\gamma}_\xi(y) = \hat{\zeta}_\xi^2(y)$ for $|y| \leq R$, where $\hat{\zeta}_\xi$ is positive definite uni-

formly continuous and has a uniformly continuous extension on $\mathbf{K}^\alpha$. Therefore, for each $c > 0$ there exists $r > 0$ such that $\|\nu_\xi - \kappa_\xi * \kappa_\xi\| < c$ for each $|\xi| > r$, where $\kappa_\xi(dx) = \zeta_\xi(x)m(dx)$ is a σ -additive non-negative measure. Hence due to corollary from Proposition IV.1.3[42] there exists $r > 0$ such that $\int_{\mathbf{K}^\alpha} \theta_\xi(y) \chi_e(-x(y)) \nu_j(dy) \geq 0$ for each $|j| > r$, consequently, $f_\xi(x) = \lim_{|j| \rightarrow \infty} \int_{\mathbf{K}^\alpha} \theta_\xi(y) \chi_e(-x(y)) \nu_j(dy) \geq 0$. From the equality $F[F(\gamma_\xi)(-y)](x) = \gamma_\xi(x)$ and the Fubini theorem it follows that

$$\int f_\xi \chi_e(y(x)) h_j(x) m(dx) = \int \theta_\xi(u + y) \nu_j(du).$$

For $y = 0$ we get

$$\lim_{|\xi| \rightarrow \infty} \int f_\xi(x) m(dx) = \int f(x) m(dx) = \lim_{|\xi| \rightarrow \infty} \lim_{|j| \rightarrow \infty} \int f_\xi(x) h_j(x) m(dx)$$

$$\text{and } \lim_{|\xi| \rightarrow \infty} \lim_{|j| \rightarrow \infty} \left| \int_{\mathbf{K}^\alpha} \theta_\xi(u) \nu_j(du) \right| \leq 1.$$

From Lemma 2.8 it follows that $\hat{f}(y) = \theta(y)$, since by Theorem 2.18 $\theta = \lim_{|\xi| \rightarrow \infty} \theta_\xi$ is a characteristic function of a probability measure on $Bf(\mathbf{K}^\alpha)$, where $f(x) = \int_{\mathbf{K}^\alpha} \theta(y) \chi_e(-x(y)) m(dy)$.

(III). Now let $\alpha = \omega_0$. It remains to show that the conditions imposed on θ are sufficient, because their necessity follows from the modification of 2.6 (since X has an algebraic embedding into X^a). The space X^a is isomorphic with $\mathbf{K}^\Lambda$ which is the space of all $\mathbf{K}$ -valued functions defined on the Hamel basis Λ in X . The Hamel basis exists due to the Kuratowski-Zorn lemma (that is, each finite system of vectors in Λ is linearly independent over $\mathbf{K}$, each vector in X is a finite linear combination over $\mathbf{K}$ of elements from Λ). Let J be a family of all non-void subsets in Λ . For each $A \in J$ there exists a functional $\theta_A : \mathbf{K}^A \rightarrow \mathbf{C}$ such that $\theta_A(t) = \theta(\sum_{y \in A} t(y)y)$ for $t \in \mathbf{K}^A$. From the conditions imposed on θ it follows that $\theta_A(0) = 1$, θ_A is uniformly continuous and bounded on $\mathbf{K}^A$, moreover, it is positive definite (or due to 2.6(6) for each $c > 0$ there are n and $q > 0$ such that for each $j > n$ and $z \in \mathbf{K}^A$ the following inequality is satisfied:

$$(i) \quad |\theta_A(z) - \theta_j(z)| \leq cbq,$$

moreover, $L_j \supset \mathbf{K}^A$, q is independent on j , c and b . From (I,II) it follows that on $Bf(\mathbf{K}^A)$ there exists a probability measure μ_A such that $\hat{\mu}_A = \theta_A$.

The family of measures $\{\mu_A : A \in J\}$ is consistent and bounded, since $\mu_A = \mu_E \circ (P_E^A)^{-1}$, if $A \subset E$, where $P_E^A : \mathbf{K}^E \rightarrow \mathbf{K}^A$ are the natural projectors. Indeed, in the case of measures with values in $\mathbf{R}$ each μ_A is the probability measure.

In view of Theorem 1.1.4[8] on a cylindrical σ -algebra of the space $\mathbf{K}^\Lambda$ there exists the unique measure μ such that $\mu_A = \mu \circ (P^A)^{-1}$ for each $A \in J$, where $P^A : \mathbf{K}^\Lambda \rightarrow \mathbf{K}^A$ are the natural projectors. From $X^a = \mathbf{K}^\Lambda$ it follows that μ is defined on $\hat{C}(X^a, X)$. For μ on $\hat{C}(X^a, X)$ there exists its extension on $Af(X, \mu)$ (see §2.1).

2.28. Definition. [36] A continuous linear operator $T : X \rightarrow Y$ for Banach spaces X and Y over $\mathbf{K}$ is called compact, if $T(B(X, 0, 1)) =: S$ is a compactoid, that is, for each neighbourhood $U \ni 0$ in Y there exists a finite subset $A \subset Y$ such that $S \subset U + co(A)$, where $co(A)$ is the least $\mathbf{K}$ -absolutely convex subset in V containing A (that is, for each a and $b \in \mathbf{K}$ with $|a| \leq 1$, $|b| \leq 1$ and for each $x, y \in V$ the following inclusion $ax + by \in V$ is accomplished).

2.29. Let B_+ be a subset of non-negative functions which are $Bf(X)$ -measurable and let C_+ be its subset of non-negative cylindrical functions. By $\hat{B}_+$ we denote a family of functions $f \in B_+$ such that $f(x) = \lim_n g_n(x)$, $g_n \in C_+$, $g_n \geq f$. For $f \in \hat{B}_+$ let

$$\int_X f(x) \mu_*(dx) := \inf_{g \geq f, g \in C_+} \int_X g(x) \mu_*(dx).$$

2.30. Lemma. A sequence of a weak distribution (μ_{L_n}) of probability Radon measures is generated by a real probability measure μ on $Bf(X)$ of a Banach space X over $\mathbf{K}$ if and only if there exists

$$(i) \quad \lim_{|\xi| \rightarrow \infty} \int_X G_\xi(x) \mu_*(dx) = 1,$$

where $\int_X G_\xi(x) \mu_*(dx) := S_\xi(\{\mu_{L_n} : n\})$ and $S_\xi(\{\mu_{L_n}\}) := \lim_{n \rightarrow \infty} \int_{L_n} F_n(\gamma_{\xi,n})(x) \mu_{L_n}(dx)$, $\gamma_{\xi,n}(y) := \prod_{l=1}^{m(n)} \gamma_\xi(y_l)$, F_n is a Fourier transformation by $(y_1, \dots, y_n)$, $y = (y_j : j \in \mathbf{N})$, $y_j \in \mathbf{K}$, $\gamma_\xi(y_l)$ are the same as in Lemma 2.8 for $\mathbf{K}^1$; here $m(n) = \dim_{\mathbf{K}} L_n < \aleph_0$, $cl(\cup_n L_n) = X = c_0(\omega_0, K)$.

Proof. If a sequence of weak distributions is generated by a measure μ , then in view of Conditions 2.6(3-6), Lemmas 2.3, 2.5, 2.8, Propositions 2.10

and 2.16, Corollary 2.13, the Lebesgue convergence theorem and the Fubini theorem, also from the proof of Theorem 2.27 and the Radon property of μ it follows that there exists $r > 0$ such that

$$\int_X G_\xi(x) \mu_*(dx) = \int_X G_\xi(x) \mu(dx) = \lim_{n \rightarrow \infty} \int_{L_n} \gamma_{\xi,n}(y) \hat{\mu}_{L_n}(y) m_{L_n}(dy),$$

since $\lim_{j \rightarrow \infty} x_j = 0$ for each $x = (x_j : j) \in X$. In addition, $\lim_{|\xi| \rightarrow \infty} S_\xi(\{\mu_{L_n}\}) = \int_X \mu(dx) = 1$. Indeed, for each $c > 0$ and $d > 0$ there exists a compact $V_c \subset X$ with $\|\mu|_{(X \setminus V_c)}\| < c$ and there exists n_0 with $V_c \subset L_n^d$ for each $n > n_0$. Therefore, choosing suitable sequences of $c(n)$, $d(n)$, $V_{c(n)}$ and L_{j_n} we get that $[\int_{L_n} \gamma_{\xi,n}(y) \hat{\mu}_{L_n}(y) m_{L_n}(dy) : n \in \mathbf{N}]$ is a Cauchy sequence, where m_{L_n} is the real Haar measure on L_n , the latter is considered as $\mathbf{Q}_p^{\mathbf{m}(n)\mathbf{b}}$, $b = \dim_{\mathbf{Q}_p} \mathbf{K}$, $m(B(L_n, 0, 1)) = 1$. Here we use $G_\xi(x)$ for a formal expression of the limit S_ξ as the integral. Then $G_\xi(x)$ (*mod* μ) is defined evidently as a function for μ or $\{\mu_{L_n} : n\}$ with a compact support, also for μ with a support in a finite-dimensional subspace L over $\mathbf{K}$ in X . By the definition $\text{supp}(\mu_{L_n} : n)$ is compact, if there is a compact $V \subset X$ with $\text{supp}(\mu_{L_n}) \subset P_{L_n} V$ for each n . That is, Condition (i) is necessary.

On the other hand, if (i) is satisfied, then for each $c > 0$ there exists $r > 0$ such that $|\int_X G_\xi(x) \mu_*(dx) - 1| < c/2$, when $|\xi| > r$, consequently, there exists n_0 such that for each $n > n_0$ the following inequality is satisfied:

$$|1 - \int_X F_n(\gamma_{\xi,n})(x) \mu_*(dx)| \leq \| \mu|_{(L_n \cap B(X, 0, R))} \| - 1 + \sup_{|x| > R} |F_n(\gamma_{\xi,n})(x)| \| \mu_{L_n}|_{(L_n \setminus B(X, 0, R))} \|.$$

Therefore, from $\lim_{R \rightarrow \infty} \sup_{|x| > R} |F_n(\gamma_{\xi,n})(x)| = 0$ and from Lemma 2.3 the statement of Lemma 2.30 follows.

2.31. Notes and definitions. Suppose X is a locally convex space over a locally compact field $\mathbf{K}$ with non-trivial non-Archimedean valuation and X^* is a topologically adjoint space. The minimum σ -algebra with respect to which the following family $\{v^* : v^* \in X^*\}$ is measurable is called a σ -algebra of cylindrical sets. Then X is called a RS -space if on X^* there exists a topology τ such that the continuity of each positive definite function $f : X^* \rightarrow \mathbf{C}$ is necessary and sufficient for f to be a characteristic functional of a non-negative measure. Such topology is called the R -Sazonov type topology. The class of RS -spaces contains all separable locally convex spaces over $\mathbf{K}$.

For example, $l^\infty(\alpha, \mathbf{K}) = c_0(\alpha, \mathbf{K})^*$, where α is an ordinal [34]. In particular we also write $c_0(\mathbf{K}) := c_0(\omega_0, \mathbf{K})$ and $l^\infty(\mathbf{K}) := l^\infty(\omega_0, \mathbf{K})$, where ω_0 is the first countable ordinal.

Let $n_{\mathbf{K}}(l^\infty, c_0)$ denotes the weakest topology on l^∞ for which all functionals $p_x(y) := \sup_n |x_n y_n|$ are continuous, where $x = \sum_n x_n e_n \in c_0$ and $y = \sum_n y_n e_n^* \in l^\infty$, e_n is the standard base in c_0 . Such topology $n_{\mathbf{K}}(l^\infty, c_0)$ is called the normal topology. The induced topology on c_0 is denoted by $n_{\mathbf{K}}(c_0, c_0)$.

2.32. Theorem. *Let $f : l^\infty(\mathbf{K}) \rightarrow \mathbf{C}$ be a functional such that*

(i) f is positive definite,

(ii) f is continuous in the normal topology $n_{\mathbf{K}}(l^\infty, c_0)$, then f is the characteristic functional of a probability measure on $c_0(\mathbf{K})$.

Proof. The case of the topological vector space X over $\mathbf{K}$ with $\text{char}(\mathbf{K}) > 0$ and a real-valued measure μ can be proved analogously to [31] for $\text{char}(\mathbf{K}) = 0$ due to §2.6 and §§2.25-2.30 (see also the proofs in Part II for $\mathbf{K}_s$ -valued measures).

2.33. Theorem. *Let μ be a probability measure on $c_0(\mathbf{K})$, then $\hat{\mu}$ is continuous in the normal topology $n_{\mathbf{K}}(l^\infty, c_0)$ on l^∞ .*

Proof. In view of Lemma 2.3 for each $\epsilon > 0$ there exists $S(\epsilon) \in c_0$ such that $\|\mu|_{L(0, S(\epsilon))}\| \geq 1 - \epsilon$, where $L(y, z) := \{x \in c_0 : |x_n - y_n| \leq |z_n|, \text{ for each } n \in \mathbf{N}\}$. Therefore,

$$|1 - \hat{\mu}(x)| \leq \epsilon + \|2\pi\eta(\xi x)\|_{C^0(L(0, S(\epsilon)))} \|\mu|_{L(0, S(\epsilon))}\|,$$

hence there exists a constant $C > 0$ such that $|1 - \hat{\mu}| \leq \epsilon + Cp_{S(\epsilon)}(x)$.

2.34. Corollary. *The normal topology $n_{\mathbf{K}}(l^\infty, c_0)$ is the R-Sazonov type topology on $l^\infty(\mathbf{K})$.*

2.35. Theorem. Non-Archimedean analog of the Minlos-Sazonov theorem. *For a separable Banach space X over $\mathbf{K}$ the following two conditions are equivalent:*

(I) $\theta : X \rightarrow \mathbf{T}$ satisfies conditions 2.6(3, 4, 5) and

for each $c > 0$ there exists a compact operator $S_c : X \rightarrow X$ such that $|\text{Re}(\theta(y) - \theta(x))| < c$ for $|\tilde{z}(S_c z)| < 1$;

(II) θ is a characteristic functional of a probability Radon measure μ

on E , where $\tilde{z}$ is an element $z \in X \hookrightarrow X^*$ considered as an element of X^* under the natural embedding associated with the standard base of $c_0(\omega_0, \mathbf{K})$, $z = x - y$, x and y are arbitrary elements of X .

Proof. ($II \rightarrow I$). For a positive definite function θ generated by a probability measure μ in view of the inequality $|\theta(y) - \theta(x)|^2 \leq 2\theta(0)(\theta(0) - \operatorname{Re}(\theta(y - x)))$ (see Propositions IV.1.1(c)[42]) and using the normalization of a measure μ by 1 we consider the case $y = 0$. For each $r > 0$ we have: $|\operatorname{Re}(\theta(0) - \theta(x))| = \int_X (1 - [\chi_e(x(u)) + \chi_e(-x(u))]/2) \mu(du) \leq \int_{B(X,0,r)} 2[\chi_e^{1/2}(x(u)) - \chi_e^{1/2}(-x(u))/(2i)]^2 \mu(du) + 2 \int_{X \setminus B(X,0,r)} \mu(du) \leq 2\pi^2 \int_{B(X,0,r)} \eta(x(u))^2 \mu(dx) + 2\mu([x : \|x\| > r])$. In view of the Radon property of the space X and Lemma 2.5 for each $b > 0$ and $\delta > 0$ there are a finite-dimensional over $\mathbf{K}$ subspace L in X and a compact subset $W \subset X$ such that $W \subset L^\delta$, $\|\mu|_{(X \setminus W)}\| < b$, hence $\|\mu|_{(X \setminus L^\delta)}\| < b$.

We consider the following expression:

$$J(j, l) := 2\pi^2 \int_{B(X,0,r)} \eta(e_j(u)) \eta(e_l(u)) \mu(du),$$

where (e_j) is the orthonormal basis in X which contains the orthonormal basis of $L = \mathbf{K}^n$, $n = \dim_{\mathbf{K}} L$. Then we choose sequences $b_j = p^{-j}$ and $0 < \delta_j < b_j$, subspaces L_j and $r = r_j$ such that $b_j r_j < 1$, $W_j \subset B(X, 0, r_j)$, $0 < r_j < r_{j+1} < \infty$ for each $j \in \mathbf{N}$ and the orthonormal basis (e_j) corresponding to the sequence $L_j \subset L_{j+1} \subset \dots \subset X$. We get, due to finiteness of $n_j := \dim_{\mathbf{K}} L_j$, that $\lim_{j+l \rightarrow \infty} J(j, l) = 0$, since $\|\mu|_{\{x: \|x\| > r_j\}}\| < b_j$, $\eta(x(u)) = 0$ for $x \in X \ominus L_j$ with $\|x\| < b_j$, $u \in B(X, 0, r_j)$. Then we define $g_{j,l} := \min\{d : d \in \Gamma_{\mathbf{K}} \text{ and } d \geq |J(j, l)|\}$, evidently, $g_{j,l} \leq p|J(j, l)|$ and there are $\xi_{j,l} \in \mathbf{K}$ with $|\xi_{j,l}|_{\mathbf{K}} = g_{j,l}$. Consequently, the family $(\xi_{j,l})$ determines a compact operator $S : X \rightarrow X$ with $\tilde{e}_j(S e_l) = \xi_{j,l}$ due to Theorem 1.2[36], where $t = \operatorname{const} \in \mathbf{K}$, $t \neq 0$. Therefore, $|\operatorname{Re}(\theta(0) - \theta(z))| < c/2 + |\tilde{z}(Sz)| < c$, $|\theta(0) - \theta(z)| < c/2 + |\tilde{z}(Sz)| < c$. We choose r such that $\|\mu|_{(X \setminus B(X,0,r))}\| < c/2$ with S corresponding to $(r_j : j)$, where $r_1 = r$, $L_1 = L$, then we take $t \in \mathbf{K}$ with $|t|c = 2$.

($I \rightarrow II$). Without restriction of generality we may take $\theta(0) = 1$ after renormalization of non-trivial θ . In view of Theorem 2.32 as in §2.6 we construct using $\theta(z)$ a consistent family of finite-dimensional distributions $\{\mu_{L_n}\}$. Let m_{L_n} be a real Haar measure on L_n which is considered as $\mathbf{Q}_{\mathbf{p}}^a$ with $a = \dim_{\mathbf{K}} L_n \dim_{\mathbf{Q}_{\mathbf{p}}} \mathbf{K}$, $m(B(L_n, 0, 1)) = 1$. In view of Proposition 2.7 and

Lemmas 2.8, 2.30: $\int_{L_n} G_\xi(x) \mu_{L_n}(dx) = \int_{L_n} \gamma_{\xi,n}(z) \theta(z) m_{L_n}(dz)$, consequently,

$$1 - \int_{L_n} F_n(\gamma_{\xi,n})(x) \mu_{L_n}(dx) = \int \gamma_{\xi,n}(z) (1 - \theta(z)) m_{L_n}(dz) =: I_n(\xi).$$

There exists an orthonormal basis in X in which S_c can be reduced to the following form $S_c = SC\hat{S}_cE$ (see Appendix), where $\hat{S}_c = \text{diag}(s_j : j \in \mathbf{N})$ in the orthonormal basis $(f_j : j)$ in X and S transposes a finite number of vectors in the orthonormal basis. That is, $|\tilde{z}(\hat{S}_c z)| = \max_j |s_j| \times |z_j|^2$. In the orthonormal basis $(e_j : j)$ adopted to $(L_n : n)$ we have $|\tilde{z}(S_c z)| = \max_{j,l \in \mathbf{N}} (|s_{j,l}| \times |z_j| \times |z_l|)$, $\|S_c\| = \max_{j,l} |s_{j,l}|$, where $S_c = (s_{j,l} : j, l \in \mathbf{N})$ in the orthonormal basis (e_j) , $r = \text{const} > 0$. In addition, $p^{-1}|x|_K \leq |x|_p \leq p|x|_K$ for each $x \in \mathbf{K}$. If S_c is a compact operator such that $|Re(\theta(y) - \theta(x))| < c$ for $|\tilde{z}(S_c z)| < 1$, $z = x - y$, then $|Re(1 - \theta(x))| < c + 2|\tilde{x}(S_c x)|$ and

$$I_n(\xi) \leq \int \gamma_{\xi,n}(z) [c + 2|\tilde{z}(S_c z)|_K] m_{L_n}(dz) \leq c + b\|S_c\|/|\xi|^2,$$

$b = \text{const}$ is independent from n , ξ and S_c ,

$$b := p \times \sup_{|\xi| > r} |\xi|^2 \int_{L_n} \gamma_{\xi,n}(z) |z|_p^2 m_{L_n}(dz) < \infty.$$

Due to the formula of changing variables in integrals (A.7[35]) the following equality is valid: $J_n(\xi) = I_n(\xi) J_n(1) / [I_n(1) |\xi|^2]$ for $|\xi| \neq 0$, where

$$J_n(\xi) = \int_{L_n} \gamma_{\xi,n}(z) |z|_p^2 m_{L_n}(dz).$$

Therefore,

$$1 - \int_X G_\xi(x) \mu_*(dx) \leq c + b\|S_c\|/|\xi|^2.$$

Then taking the limit with $|\xi| \rightarrow \infty$ and then with $c \rightarrow +0$ with the help of Lemma 2.30 we get the statement $(I \rightarrow II)$.

2.36. Definition. Let on a completely regular space X with $\text{ind}(X) = 0$ two non-zero real-valued measures μ and ν are given. Then ν is called absolutely continuous relative to μ if $\nu(A) = 0$ for each $A \in Bf(X)$ with $\mu(A) = 0$ and it is denoted $\nu \ll \mu$. Measures ν and μ are singular to each other if there is $F \in Bf(X)$ with $|\mu|(X \setminus F) = 0$ and $|\nu|(F) = 0$ and it is denoted $\nu \perp \mu$. If $\nu \ll \mu$ and $\mu \ll \nu$ then they are called equivalent, $\nu \sim \mu$.

2.37. For a Banach space X over $\mathbf{K}$ and σ -algebra $\mathbf{B} \supset Bf(X)$ with a real probability measure μ and a σ -subalgebra $\mathbf{B}_0 \subset \mathbf{B}$ analogously to §13[38] a function $\bar{\mu}(A|x)$ satisfying three conditions:

- (a) $\bar{\mu}(A|x)$ is $\mathbf{B}_0$ -measurable by x for each $A \in \mathbf{B}$;
- (b) $\mu(A \cap A_0) = \int_{A_0} \bar{\mu}(A|x) \mu(dx) =: \mu_A(A_0)$ for each $A_0 \in \mathbf{B}_0$;
- (c) $\bar{\mu}(A|x)$ is a measure by $A \in \mathbf{B}$ for almost all x relative to a measure μ , then it is called a conditional measure corresponding to μ relative to a σ -algebra $\mathbf{B}_0$.

Then we define $\mu(A|x)$ by formula (b) and we then redefine it on the set of measure zero relative to μ for each A . Let $\theta(z|x) = \theta_1(z|x) + i\theta_2(z|x)$ with $\int_{A_0} [\chi_e(\tilde{z}(x)) + \chi_e(-\tilde{z}(x))]/2 \mu(dx) = \int_{A_0} \theta_1(z|x) \mu(dx)$; $\int_{A_0} [\chi_e(\tilde{z}(x)) - \chi_e(-\tilde{z}(x))]/(2i) \mu(dx) = \int_{A_0} \theta_2(z|x) \mu(dx)$, where θ_j are measurable functions, $z \in X$, $\tilde{z} \in X'$ corresponds to z under the natural embedding of X into X' , $x \in X$, $A_0 \in \mathbf{B}_0$. Then we choose analogously to [38] sets $A_{n,k}$ with $\text{diam}(A_{n,k}) < p^{-n}$, using §§2.7, 2.8, 2.30 and the proof of §2.27, substituting $\exp(-(x, x)/n)$ on $G_\xi(x)$, defining sets B_n , G , G_1 , we get for each $c > 0$ that there exists $r > 0$ such that

$$\begin{aligned} \text{Re}(1 - \theta(z|x)) &\leq \int_X G_\xi(y) (1 - [\chi_e(\tilde{z}(y)) + \chi_e(\tilde{z}(x))]/2) \mu(dy|x) + 2 \int_X (1 - G_\xi(y)) \mu(dy|x) \\ &\leq \int (2\pi\eta^2(\tilde{z}(y)) G_\xi(y) \mu(dy|x) / 2 + c/2 \end{aligned}$$

for each $|\xi| > r$ and $x \in G_1 \cap G$, since from $\|z\| \times \|y\| \leq 1$ it follows that $(1 - \chi_e(\tilde{z}(y))) = 0$ and $|1 - \exp(it)| \leq |t|$ for $t \in \mathbf{R}$, $i = (-1)^{1/2}$.

For $C(z_1, z_2)(x) := \int_X \eta(\tilde{z}_1(y)) \eta(\tilde{z}_2(y)) G_\xi(y) \mu(dy|x)$ we have $C(0, 0) = 0$ and $C(z, z)(x) = \int_{X \setminus B(X, 0, r)} \eta^2(\tilde{z}(y)) G_\xi(y) \mu(dy|x)$ for $\|z\| = 1/r > 0$. Since $\theta(z|x) = \int_X \chi_e(\tilde{z}(y)) \mu(dy|x)$, considering projectors $P_{L_n} : X \rightarrow L_n$ onto finite-dimensional subspaces over $\mathbf{K}$, $L \subset X$, we get $\theta_L(z|x)$ corresponding to $\mu_L(dx)$ and $\mu_L(dy|x)$. At the same time θ_L satisfies Conditions 2.6(3,4,5), consequently, $\mu_L(A|x)$ are σ -additive by A for fixed x , since $\theta(z|x)$ are continuous by z for $x \in G \cap G_1$. From the Prohorov theorem IX.4.2.1[5], Lemma 2.5, $x \in G \cap G_1$ and $C(e_j, e_l)(x) = \int_{L_{j,l} \setminus B(L_{j,l}, 0, 1)} \eta(y_l) \eta(y_j) F_2(\gamma_{\xi, 2})(y_j, y_l) \mu_{L_{j,l}}(dy_j dy_l|x)$ it follows that $\lim_{j+l \rightarrow \infty} \int_{L_{j,l}} \psi(x) C(e_j, e_l)(x) \mu_{L_{j,l}}(dx) = \lim_{j+l \rightarrow \infty} \int_{L_{j,l}} \psi(x) \int_{L_{j,l}} \eta(y_j) \eta(y_l) \chi_e(\tilde{z}(y)) \mu_{L_{j,l}}(dy|x) \mu_{L_{j,l}}(dx) = 0$ for each bounded $\mathbf{B}_0$ -measurable function $\psi(x)$, whence $\lim_{n \rightarrow \infty} \mu([x : |C(e_j, e_l)(x)| > c \text{ for some } j + l > n]) = 0$ for each $c > 0$, where $L_{j,l} = Ke_j \oplus Ke_l$, $y_j \in \mathbf{K}$.

Let $G_2 := [x : \text{for each } c > 0 \text{ there exists } n > 0 \text{ with } |C(e_j, e_l)(x)| < c \text{ for each } j + l > n] \subset X \setminus (\bigcup_{c>0} \bigcap_{n=1}^{\infty} \bigcup_{j+l>n} [x : |C(e_j, e_l)(x)| > c])$, then $\mu(G_2) = 1$. Therefore, for each $x \in G \cap G_1 \cap G_2$ there exists a compact operator $D : X \rightarrow X$ such that $|C(z_1, z_2)(x)| \leq |\tilde{z}_2(Dz_1)|$ for each $z_1, z_2 \in X$. Using $\chi_e(\tilde{z}(x))$ (instead of $\exp(2\pi i(z, x))$ from [38]) and Theorem 2.35 we find a conditional measure $\bar{\mu}(A|x)$.

2.38. Remark. For $\mu : Bf(X) \rightarrow \mathbf{R}$ martingales, semi-martingales and sub-martingales are defined analogously §14[38]. Then in the latter case proofs of Lemmas 1,2 and Theorems 1,2 differ slightly (with substitution of a Hilbert space on a Banach space X over $\mathbf{K}$ in all formulations).

3 Quasi-invariant measures.

In this section after few preliminary statements there are given the definition of a quasi-invariant measure and the theorems about quasi-invariance of measures relative to transformations of a Banach space X over $\mathbf{K}$.

3.1. Let X be a Banach space over $\mathbf{K}$, $(L_n : n)$ be a sequence of subspaces, $cl(\bigcup_n L_n) = X$, $L_n \subset L_{n+1}$ for each n , μ^j be probability measures, $\mu^2 \ll \mu^1$, $(\mu_{L_n}^j)$ be sequences of weak distributions, also let there exist derivatives $\rho_n(x) = \mu_{L_n}^2(dx)/\mu_{L_n}^1(dx)$ and the following limit $\rho(x) := \lim_{n \rightarrow \infty} \rho_n(x)$ exists.

Theorem. *If μ^j are real-valued and in addition $\int_X \rho(x) \mu^1(dx) = 1$ with $\rho \in L^1(\mu^1)$, then this is equivalent to the following: there exists $\rho(x) = \mu^2(dx)/\mu^1(dx) \pmod{\mu^1}$.*

Proof may be obtained from the proof of theorem 1 §15[38] by simple modification substituting a Hilbert space over $\mathbf{R}$ onto the Banach space X over $\mathbf{K}$ (see also §2 above).

3.2. Theorem. *Measures $\mu^j : Bf(X) \rightarrow \mathbf{R}$, $j = 1, 2$, for a Banach space X over $\mathbf{K}$ are orthogonal $\mu^1 \perp \mu^2$ if and only if $\rho(x) = 0 \pmod{\mu^1}$.*

Proof. In the case of real-valued μ^j the proof differs only slightly from the proof of Theorem 2 §15[38] (see also the proof in Part II for $\mathbf{K}_s$ -valued measures).

3.3. Note. For real-valued measures μ^j on $Bf(X)$ for a Banach space X over $\mathbf{K}$ (instead of a Hilbert space) using the above given statements Theorems 3-6 and Corollary in §15[38] may be reformulated and proved. For real-valued μ_j see also Kakutani Theorem 4.1 §II.4.6[8].

3.4. Theorem. Let ν, μ, ν_j, μ_j be probability measures with values in $\mathbf{R}$, $X = \prod_{j=1}^{\infty} X_j$ be a product of completely regular spaces X_j with $\text{ind}(X_j) = 0$ and $\mu \ll \nu$, then $\mu_j \ll \nu_j$ for each j and $\prod_{j=1}^{\infty} \beta_j$ converges to β , $\infty > \beta > 0$, where $\beta_j := \|(\rho_j)^{1/2}\|_{L^1(X_j, \nu_j)}$, $\rho_j(x) = \mu_j(dx)/\nu_j(dx)$.

Proof. The case of measures with values in $\mathbf{R}$ may be obtained by a modification of Theorem 1 §16[38] (see also the proof in Part II for $\mathbf{K}_s$ -valued measures).

3.5. Definition. Let X be a Banach space over $\mathbf{K}$, Y be a completely regular space with $\text{ind}(X) = 0$, $\nu : Bf(Y) \rightarrow \mathbf{R}$, $\mu^y : Bf(X) \rightarrow \mathbf{R}$ for each $y \in Y$, suppose $\mu^y(A) \in L^1(Y, \nu, \mathbf{R})$ for each $A \in Bf(X)$. Then we define:

$$(i) \mu(A) = \int_Y \mu_y(A) \nu(dy).$$

A measure μ is called mixed. We define measures π^j by the formula:

$$(ii) \pi^j(A \times C) = \int_C \mu^{j,y}(A) \nu^j(dy),$$

where $j = 1, 2$ and $\mu^{j,y}$ together with ν^j are defined as above μ^y and ν . For μ^y and ν with values in $\mathbf{R}$ definitions of μ and π^j , Theorems 1,2 and Lemma 1 §18[38] may be modified for the non-Archimedean case of the Banach space X over $\mathbf{K}$.

3.6. Definition. For a Banach space X over $\mathbf{K}$ an element $a \in X$ is called an admissible shift of a measure μ , if $\mu_a \ll \mu$, where $\mu_a(A) = \mu(S_{-a}A)$ for each A in $Bf(X)$, $S_a A := a + A$, $\rho(a, x) := \rho_\mu(a, x) := \mu_a(dx)/\mu(dx)$, $M_\mu := [a \in X : \mu_a \ll \mu]$ (see §§2.1 and 2.36).

The proofs of the properties given below in §3.7 differ slightly from §19[38], when a Hilbert space is substituted on a Banach space X over $\mathbf{K}$ (see the proofs in Part II for $\mathbf{K}_s$).

3.7. Properties of M_μ and ρ from §3.6.

I. The set M_μ is a semigroup by addition, $\rho(a + b, x) = \rho(a, x)\rho(b, x - a)$ for each $a, b \in M_\mu$.

II. If $a \in M_\mu$, $\rho(a, x) \neq 0 \pmod{\mu}$, then $\mu_a \sim \mu$, $-a \in M_\mu$ and $\rho(-a, x) = 1/\rho(a, x - a)$.

III. If $\nu \ll \mu$ and $\nu(dx)/\mu(dx) = g(x)$, then $M_\mu \cap M_\nu = M_\mu \cap [a : \mu([x : g(x) = 0, g(x - a)\rho_\mu(a, x) \neq 0]) = 0]$ and $\rho_\nu(a, x) = [g(x - a)/g(x)]\rho_\mu(a, x) \pmod{\nu}$ for $a \in M_\mu \cap M_\nu$.

IV. If $\nu \sim \mu$, then $M_\nu = M_\mu$.

V. Also (V-VIII) §19[38] may be slightly reformulated and proved for real-valued measures.

3.8. Definition and notes. A linear operator U on X has a matrix representation $u(i, j) = \tilde{e}_i(Ue_j)$, where $(e_j : j)$ is a non-Archimedean orthonormal basis in X , $\tilde{e}_i$ are vectors of the topologically adjoint space X^* under the natural embedding (see §§2.2, 2.6). For a compact operator $U - I$ we get that there exists $\lim_{n \rightarrow \infty} \det\{\hat{r}_n(U)\} \in \mathbf{K}$ which we adopt as the definition of $\det\{U\}$, where $r_n : X \rightarrow \mathbf{K}^n$ is the projection operator with $\hat{r}_n(U) := \{u(i, j) : i, j = 1, \dots, n\}$ (see also Appendix, use that $\lim_{i+j \rightarrow \infty} u(i, j) = 0$).

3.9. Theorem. Let X be a Banach space over $\mathbf{K}$, μ be a probability real-valued measure and $T : X \rightarrow X$ be a compact operator such that $\operatorname{Re}(1 - \hat{\mu}(z)) \rightarrow 0$ for $|\tilde{z}(T^2 z)| \rightarrow 0$ and $z \in X$, $\tilde{z} \in X'$ corresponding to z . Then $M_\mu \subset (TX)^\sim$, where $Y^\sim$ is a completion by $\|\cdot\|_Y$ of a normed space Y .

Proof. For each $a \in M_\mu$ and $b > 0$ we have $J(z) = \int_X [1 - [\chi_e(\tilde{z}(x)) + \chi_e(-\tilde{z}(x))]/2] \rho(a, x) \mu(dx) \leq J_1(z) + J_2(z)$, where $J_1(z) := b \int_X [1 - [\chi_e(\tilde{z}(x)) + \chi_e(-\tilde{z}(x))]/2] \mu(dx)$, $J_2(z) = 2 \int_{\{x: \rho(a, x) > b\}} \mu(dx)$, $J_1(z) \rightarrow 0$ for $|\tilde{z}(T^2 z)| \rightarrow 0$, $J_2(z) \rightarrow 0$ for $b \rightarrow +\infty$, consequently, $\lim_{|\tilde{z}(T^2 z)| \rightarrow 0} \operatorname{Re}(\mu_a(X) - \chi_e(\tilde{z}(a)) \hat{\mu}(z)) = 0$. From $|1 - \hat{\mu}(z)|^2 \leq 2 \operatorname{Re}(1 - \hat{\mu}(z))$ and $\mu(X) = \mu_a(X) = 1$ it follows that $\chi_e(\tilde{z}(a)) \rightarrow 1$ and $\tilde{z}(a) \rightarrow 0$ for $|\tilde{z}(T^2 z)| \rightarrow 0$. If $|\tilde{z}(a)| < \epsilon$ for $|\tilde{z}(T^2 z)| < \delta$, then $|\tilde{z}(a)|^2 < \epsilon^2 |\tilde{z}(T^2 z)| / \delta$. Let $(e_j : j)$ be orthonormal basis in X such that $T = (T_{i,j} : i, j \in \mathbf{N})$, $\lim_{i+j \rightarrow \infty} T_{i,j} = 0$, we denote $D_j := \sup_k |T_{k,j}|$, hence $|a_j|^2 < \epsilon^2 D_j^2 / \delta$, $a \in (TX)^\sim$, $\|a\|_{TX} := \sup_j D_j |a_j| < \infty$ (see Appendix).

3.10. Corollary. Let X and μ be the same as in §3.9, then $L \cap M_\mu$ is a set of the first category in X for each infinite-dimensional linear subspace $L \subset X$ over $\mathbf{K}$.

Proof follows from the fact that $(TB(X, 0, p^j))^\sim =: A_j$ is nowhere dense in $L \cap B(X, 0, p^j)$ for each $j \in \mathbf{N}$, since A_j is compact in X [36].

The proof of Theorem 2 §19[38] can not be transferred on non-Archimedean X , there exists a compact V in $X = c_0(\omega_0, \mathbf{K})$ and its linear span $Y = sp_{\mathbf{K}} V$ such that an ultranorm $p_E(x) := \inf\{|a| : x \in aE\}$ (see Exer. 6.204 and 5.202[33]) produces from Y non-separable and non-Radonian Banach space $l^\infty(\omega_0, \mathbf{K})$ [8, 34], where $E = co(V)$ in Y .

Note. Moreover, as it will be seen below from the proofs of Theorems 3.20 and 4.2 it follows the existence of a probability quasi-invariant measure $\mu : Bf(X) \rightarrow \mathbf{R}$ with $M_\mu \supset H + G_T$, where G_T is a compact subgroup in X with $\mu(G_T) > 0$ such that $\mu(L) = 0$ for real-valued μ and each linear

subspace L in X with $\dim_{\mathbf{K}} L < \aleph_0$ (see Part II also respectively).

3.11. Definition. For a Banach space X over $\mathbf{K}$ and a measure $\mu : Bf(X) \rightarrow \mathbf{R}$, $a \in X$, $\|a\| = 1$, a vector a is called an admissible direction, if $a \in M_\mu^{\mathbf{K}} := [z : \|z\|_X = 1, \lambda z \in M_\mu \text{ and } \rho(\lambda z, x) \neq 0 \pmod{\mu} \text{ (relative to } x)]$ and for each $\lambda \in \mathbf{K}] \subset X$. Let $a \in M_\mu^{\mathbf{K}}$ we denote by $L_1 := \mathbf{K}a$, $X_1 = X \ominus L_1$, μ^1 and $\tilde{\mu}^1$ are the projections of μ onto L_1 and X_1 respectively, $\tilde{\mu} = \mu^1 \otimes \tilde{\mu}^1$ be a measure on $Bf(X)$, given by the the following equation $\tilde{\mu}(A \times C) = \mu^1(A) \tilde{\mu}^1(C)$ on $Bf(L_1) \times Bf(X_1)$ and extended on $Bf(X)$, where $A \in Bf(L_1)$, $C \in Bf(X_1)$.

3.12. Theorem. For a to be in $M_\mu^{\mathbf{K}}$ with a real-valued probability measure μ it is necessary and sufficient that the following conditions be satisfied (i-iv): (i) $\mu \ll \tilde{\mu}$; (ii) $\mu^1 \ll m$, where m is a real-valued Haar measure on L_1 ; (iii) $\mu^1(dx)/\mu(dx) := h(x) \neq 0$ for $|a(x)| > 0$; (iv) $\mu([x : g(x - \lambda a) \neq 0, g(x) = 0]) = 0$ for $g(x) = \mu(dx)/\tilde{\mu}(dx)$ whilst $|\lambda| > 0$.

Proof. From $\tilde{\mu}_{\lambda a} = \mu_{\lambda a}^1 \otimes \tilde{\mu}^1$ and §§3.10 II,III sufficiency follows. The proof of necessity is analogous to that of Theorem 1 §20[38] (see §3.7). Indeed, from $\rho(\lambda z, x) \neq 0 \pmod{\mu}$ it follows that $\mu^1 \sim m$ due to Proposition 11 §VII.1.9[5].

3.13. Note. For $\mathbf{K}_s$ -valued measures Theorem 3.12 is untrue, since there are cases with μ that are not absolutely continuous relative to $\tilde{\mu}$. For example, for a probability measure $\mu = \mu_1 + \mu_2 \neq 0$, $\mu_2(X) = 0$ with μ_1^1 equivalent to the Haar measure m on L_1 , $\mu_2^1 = 0$, when for μ_2 all atoms $(a_{j,2})$ are points and for each $x \in L_1$ is accomplished $\sum_{P_{L_1}(a_{j,2})=x} \mu_2(a_{j,2}) = 0$ and atoms $(a_{j,1})$ of μ_1 are such that $P_{L_1}(a_{j,1}) \neq P_{L_1}(a_{l,2})$ for each j, l . We can choose $(a_{j,s})$ such that there exists a compact $S \subset X$ with $\mu_1(S) = 0$, $\mu_2(S) \neq 0$, but $\tilde{\mu}(S) = 0$, since $\mu_2^1 = 0$.

3.14. Definition and notes. A measure $\mu : Bf(X) \rightarrow \mathbf{R}$ for a Banach space X over $\mathbf{K}$ is called a quasi-invariant measure if M_μ contains a $\mathbf{K}$ -linear manifold J_μ dense in X .

From §3.7 and Definition 3.11 it follows that $J_\mu \subset M_\mu^{\mathbf{K}}$.

Let $(e_j : j \in \mathbf{N})$ be an orthonormal basis in X , $H = sp_{\mathbf{K}}(e_j : j)$. We denote $\Omega(\mathbf{R}) = [\mu | \mu \text{ is a real measure with a finite total variation on } Bf(X) \text{ and } H \subset J_\mu]$.

A measure $\mu \in \Omega(\mathbf{R})$ that can not be represented as a sum of two singular to each other measures from $\Omega(\mathbf{R})$ is called an extremal measure. $Bf(X)$ -measurable function $h : X \rightarrow \mathbf{R}$ is called invariant for $\mu \in \Omega(\mathbf{R})$, if $h(a+x) =$

$h(x)$ μ -a.e. by x for each $a \in H$.

3.15. Lemma. *If $\mu \in \Omega(\mathbf{R})$ and h is a bounded invariant function for μ , then $h(x) = \int h(y)\mu(dy, \mathbf{B}^n|x)$ μ -a.e. for each n , where $\mathbf{B}^n = P_{X^n}^{-1}(Bf(X^n))$, $H_n = sp_K(e_1, \dots, e_n)$, $X^n = X \ominus H_n$, $\mathbf{B}_n = P_{H_n}^{-1}(Bf(H_n))$, $P_{X^n} : X \rightarrow X^n$, $P_{H_n} : X \rightarrow H_n$ is projector $\mu(*, \mathbf{B}^n|x)$ is a conditional measure.*

Proof. From §2.37 the existence of a conditional measure follows. Lemmas 1, 2 and corollary 1 §23[38] may be transferred on a Banach space X over $\mathbf{K}$. Let us take $h_\xi(x) = \int_{H_n} h(x+y)\gamma_{\xi,n}(x)m_{H_n}(dx)$, where $\gamma_{\xi,n}$ corresponds to H_n and m_{H_n} are the same as in §2.30. Choosing $|\xi_l| \rightarrow \infty$ whilst $l \rightarrow \infty$ as in Lemma 3 §23[38] we get the statement of Lemma 3.15.

3.16. Theorem. *A measure $\mu \in \Omega(\mathbf{R})$ is an extremal measure if and only if μ is equivalent to $\nu \in \mathbf{R}$, where $\mathbf{R} := [\nu \in \Omega(\mathbf{R})]$ for each n there exists $m > n$ such that $\nu(A \cap B) = \nu(A)\nu(B)$ for each $A \in \mathbf{B}_n$, $B \in \mathbf{B}^m$.*

Proof may be obtained by a slight modification of the proof of Theorem 1 §23[38] with the help of Theorems 3.1, 3.2, 3.11, Note 3.3 and §§3.5, 2.38, since $\mu : Bf(X) \rightarrow \mathbf{R}$. In $\mathbf{R}$ the notion of the infinite number of crossings of a segment $[\beta_1, \beta_2] \subset \mathbf{R}$ by a sequence $(\alpha_j : j \in \mathbf{N})$ is kept (that is, there are $k_1 < k_2 < \dots$ such that $\alpha_{k_1} \geq \beta_2$, $\alpha_{k_2} \leq \beta_1, \dots, \alpha_{k_{2n-1}} \geq \beta_2$, $\alpha_{k_{2n}} \leq \beta_1$). Then a martingale (ϕ_n) with values in $\mathbf{R}$ may be considered, $B_{\beta_1, \beta_2} := [x \in X : (\phi_n(x) : n \in \mathbf{N}) \text{ crosses an infinite number of times a segment } (\beta_1, \beta_2) \subset \mathbf{R}]$, where $\beta_2 > \beta_1 \geq 0$, n is fixed, $\rho_n^m(x|y) = \mu_n^m(dx|y)/\mu^m(dx)$, $\mu_n^m(*|y) := \mu(*, \mathbf{B}_n|y)$ on $\mathbf{B}^m$ for $n \leq m$, $\mu^m := \mu|_{\mathbf{B}^m}$. In addition corollaries from Lemma 3 §23 [38] may be slightly modified and proved for the Banach space X over $\mathbf{K}$ due to Lemma 3.15 and Theorem 2.35. Then the Lebesgue measure on $\mathbf{R}^b$ must be substituted into the Haar measure on $\mathbf{Q}_p^b$ with values in $[0, \infty)$. [Here the situation is analogous to Theorem 1 §16[38] about absolute continuity of real-valued product-measures μ on a Hilbert space X_0 and its more general in some respects variant Theorem II.4.1[8] on products of measurable spaces. Indeed, $Bf(X_0)$ for X_0 in a topology of $\|\cdot\|$ and a weak topology $\sigma(X_0, X_0^*)$ coincide.]

3.17. Theorem. *For a Banach space X over $\mathbf{K}$ and each $\mu \in \Omega(\mathbf{R})$ there exists $[\mu^y : y \in Y] \subset \mathbf{R}$ and a measure ν on $\mathbf{B}^\infty$ such that $\mu(A) = \int_X \mu^y(A)\nu(dy)$ for each $A \in Bf(X)$, where $\mathbf{B}^\infty := (\cap_n \bar{\mathbf{B}}^n \cap Bf(X)$, $\bar{\mathbf{B}}^n$ is a completion of $\mathbf{B}^n$ by a measure μ , $\nu = \mu|_{\mathbf{B}^\infty}$, $\mu^y(A)$ is $Af(X, \mu)$ -measurable (see the notation in §§2.1, 3.14 and 3.16).*

Proof. From §2.37 for a conditional measure $\mu^y(A) := \mu(A, \mathbf{B}^\infty|y)$ corresponding to μ relative to $\mathbf{B}^\infty$ it follows that $\mu(A) = \int \mu(A, \mathbf{B}^\infty|y)\mu(dy)$.

Let m be the real-valued Haar measure on $Bf(\mathbf{K})$, $m \otimes \nu$ be a measure on $Bf(\mathbf{K}) \times \mathbf{B}^\infty$, then as in the proof of Theorem 2 §23[38] we establish that $\mu_{\tau a}^y \sim \mu^y$ for $m \otimes \nu$ -a.e. pairs $(\tau, y) \in \mathbf{K} \times Y$, $Y = X$. Therefore, for ν -a.e. y the set $S_y := [\tau : \mu_{\tau a}^y \sim \mu^y, \tau \in \mathbf{K}]$ has $m(\mathbf{K} \setminus S_y) = 0$ and $S_y \in Af(\mathbf{K}, m)$, moreover, S_y is the additive group, consequently, $\nu([y : S_y \neq \mathbf{K}]) = 0$. Then we may continue analogously to §23[38].

3.18. Theorem. *If $\mu : Bf(Y) \rightarrow \mathbf{R}$ is a σ -finite measure on $Bf(Y)$, Y is a complete separable ultrametrizable $\mathbf{K}$ -linear subspace such that $co(S)$ is nowhere dense in Y for each compact $S \subset Y$, where $\mathbf{K}$ is an infinite non-discrete non-Archimedean field with a multiplicative ultranorm $|\cdot|_{\mathbf{K}}$. Then from $J_\mu = Y$ it follows that $\mu = 0$.*

Proof. Since μ is σ -finite, then there are $(Y_j : j \in H) \subset Bf(Y)$ such that $Y = \bigcup_{j \in H} Y_j$ and $0 < \|\mu|_{Bf(Y_j)}\| \leq 1$ for each j , where $H \subset \mathbf{N}$, $Y_j \cap Y_l = \emptyset$ for each $j \neq l$. If $\text{card}(H) = \aleph_0$, then we define a function $f(x) = 1/[2^j |\mu|(Y_j)]$ for real-valued μ . Then we define a measure $\nu(A) = \int_A f(x) \mu(dx)$, $A \in Bf(Y)$. Therefore, $|\nu|(Y) \leq 1$ and $J_\nu = Y$, since $f \in L^1(Y, \mu, \mathbf{R})$. Hence it is sufficient to consider μ with $\|\mu\| \leq 1$ and $|\mu|(Y) = 1$. For each $n \in \mathbf{N}$ in view of the Radonian property of Y there exists a compact $X_n \subset Y$ such that $|\mu|(Y \setminus X_n) < 1/n$. In Y there is a countable everywhere dense subset $(x_j : j \in \mathbf{N})$, hence $Y = \bigcup_{j \in \mathbf{N}} B(Y, x_j, r_l)$ for each $r_l > 0$, where $B(Y, x, r_l) = [y \in Y : d(x, y) \leq r_l]$, d is an ultrametric in Y , i.e. $d(x, z) \leq \max(d(x, y), d(y, z))$, $d(x, z) = d(z, x)$, $d(x, x) = 0$, $d(x, y) > 0$ for $x \neq y$ for each $x, y, z \in Y$. Therefore, for each $r_l = 1/l$, $l \in \mathbf{N}$ there exists $k(l) \in \mathbf{N}$ such that $|\mu|(X_{n,l}) > 1 - 2^{-l-n}$, where $X_{n,l} := \bigcup_{j=1}^{k(l)} B(Y, x_j, r_l)$, consequently, $|\mu|(Y \setminus X_n) \leq 2^{-n}$ for $X_n := \bigcap_{l=1}^\infty X_{n,l}$. The subsets X_n are compact, since X_n are closed in Y and the metric d on X_n is completely bounded and Y is complete (see Theorems 3.1.2 and 4.3.29 [10]). Then $0 < |\mu|(X) \leq 1$ for $|\mu|(Y \setminus X) = 0$ for $X := sp_{\mathbf{K}}(\bigcup_{n=1}^\infty X_n)$.

The sets $\tilde{Y}_n = co(Y_n)$ are nowhere dense in Y for $Y_n = \bigcup_{l=1}^n X_l$, consequently, $sp_{\mathbf{K}} Y_n$ are nowhere dense in Y . Moreover, $(Y \setminus \bigcup_{n=1}^\infty Y_n) \neq \emptyset$ is dense in Y due to the Baire category theorem (see 3.9.3 and 4.3.26 [10]). Therefore, $y + X \subset Y \setminus X$ for $y \in Y \setminus X$ and from $J_\mu = Y$ it follows that $|\mu|(X) = 0$, since $|\mu|(y + X) = 0$ (see §§2.38 and 3.14 above). Hence we get the contradiction, consequently, $\mu = 0$.

3.19. Corollary. *If Y is a Banach space or a complete countably-ultranormable infinite-dimensional over $\mathbf{K}$ space, $\mu : Bf(Y) \rightarrow \mathbf{R}$ and $J_\mu =$*

Y , then $\mu = 0$.

Proof. The space Y is evidently complete and ultrametrizable, since its topology is given by a countable family of ultranorms. Moreover, $co(S)$ is nowhere dense in Y for each compact S in Y , since $co(S) = cl(S_{bc})$ is compact in Y and does not contain in itself any open subset in Y due to (5.7.5)[33] (for Y over $\mathbf{R}$ and real-valued measures see Theorem 4 §V.5.3[13]).

3.20. Theorem. *Let X be a separable Banach space over a locally compact infinite field $\mathbf{K}$ with a nontrivial valuation such that either $\mathbf{K} \supset \mathbf{Q}_p$ or $\text{char}(\mathbf{K}) = p > 0$. Then there are probability measures μ on X with values in $\mathbf{R}$ such that μ are quasi-invariant relative to a dense $\mathbf{K}$ -linear subspace J_μ .*

Proof. Let $S(j, n) := p^j B(\mathbf{K}, 0, 1) \setminus p^{j+1} B(\mathbf{K}, 0, 1)$ for $j \in \mathbf{Z}$ and $j \leq n$, $S(n, n) := p^n B(\mathbf{K}, 0, 1)$, w be the Haar measure on $\mathbf{K}$ considered as the additive group (see [15, 34]) with values in $\mathbf{R}$. Then for each $c > 0$ and $n \in \mathbf{N}$ there are measures m on $Bf(\mathbf{K})$ such that $m(dx) = f(x)v(dx)$, $|f(x)| > 0$ for each $x \in \mathbf{K}$ and $|m(p^n B(\mathbf{K}, 0, 1)) - 1| < c$, $m(\mathbf{K}) = 1$, $|m|(E) \leq 1$ for each $E \in Bf(\mathbf{K})$, where $v = w$, $v(B(\mathbf{K}, 0, 1)) = 1$. Moreover, we can choose f such that a density $m_a(dx)/m(dx) =: d(m; a, x)$ be continuous by $(a, x) \in \mathbf{K}^2$ and for each $c' > 0$, x and $|a| \leq p^{-n}$: $|d(m; a, x) - 1| < c'$. For this we can define f for $v = w$, for example, to be at the beginning locally constant such that $f(x) = a(j, n)$ for $x \in S(j, n)$, where $a(j, n) = r^{n(j-n)}(1 - r^{-n})(1 - 1/p)p^{-n}$ for $j < n$, $a(n, n) = (1 - r^{-2n})p^{-n}$ and $m(E) := \sum \{a(j, n)v(p^{n-j}(E \cap S(j, n)))/v(p^{n-j} S(j, n)) : j \in \mathbf{Z}, j \leq n\}$. Then we can take $g(x) = f(x) + h(x)$ and $y(dx) := g(x)v(dx)$, and a continuous $h(x) : K \rightarrow \mathbf{R}$, with $\sup\{|h(x)/f(x)| : x \in K\} \leq c''$ and $0 < c'' \leq 1/p^n$.

Let $\{m(j; dx)\}$ be a family of measures on $\mathbf{K}$ with the corresponding sequence $\{k(j)\}$ such that $k(j) \leq k(j+1)$ for each j and $\lim_{i \rightarrow \infty} k(i) = \infty$, where $m(j; dx)$ corresponds to the partition $[S(i, k(j))]$. The Banach space X is isomorphic with $c_0(\omega_0, \mathbf{K})$ [34]. It has the orthonormal basis $\{e_j : j = 1, 2, \dots\}$ and the projectors $P_j x = (x(1), \dots, x(j))$ onto $\mathbf{K}^j$, where $x = x(1)e_1 + x(2)e_2 + \dots$. Then there exists a cylindrical measure μ generated by a consistent family of measures $y(j, B) = b(j, E)$ for $B = P_j^{-1}E$ and $E \in Bf(\mathbf{K}^j)$ [5, 8] where $b(j, dz) = \otimes [m(j; dz(i)) : i = 1, \dots, j]$, $z = (z(1), \dots, z(j))$. Let $L := L(t, t(1), \dots, t(l); l) := \{x : x \in X \text{ and } |x(i)| \leq p^a, a = -t - t(i) \text{ for } i = 1, \dots, l, \text{ and } a = -k(j) \text{ for } j > l\}$, then L is compact in X , since X is Lindelöf and L is sequentially compact [10]. Therefore, for each $c > 0$ there exists L such that $|\mu(X \setminus L)| < c$, since there is $l \in \mathbf{N}$ with

$|1 - \prod[m(j; p^{k(j)} B(K, 0, 1)) : j > l]| < c/2$ (or due to the choice of $a(j, n)$).

In view of the Prohorov theorem §IX.4.2[5] for real measures and due to Lemma 2.3 μ has the countably-additive extension on $Bf(X)$, consequently, also on the complete σ -field $Af(X, \mu)$ and μ is the Radon measure.

Let $z' \in sp_{\mathbf{K}}\{e_j : j = 1, 2, \dots\}$ and $z'' = \{z(j) : z(j) = 0 \text{ for } j \leq l \text{ and } z(j) \in S(n, n), j = 1, 2, \dots, n = k(j)\}$, $l \in \mathbf{N}$, $z = z' + z''$. In view of the Kakutani theorem [8] and also I.1.4, II.4.1[34] there are $m(j; dz(j))$ such that $\rho_\mu(z, x) = \prod\{d(j; z(j), x(j)) : j = 1, 2, \dots\} = \mu_z(dx)/\mu(dx) \in L^1(X, \mu, \mathbf{R})$ for each such z and $x \in X$, where $d(j; *, *) = d(m(j; *), *, *)$ and $\mu_z(X) = \mu(X) = 1$.

When $\text{char}(\mathbf{K}) = 0$, then in this proof it is also possible to consider $S(j, n) := \pi^j B(\mathbf{K}, 0, 1) \setminus \pi^{j+1} B(\mathbf{K}, 0, 1)$, where $\pi \in \mathbf{K}$, $p^{-1} \leq |\pi| < 1$, π is the generator of the valuation group of $\mathbf{K}$.

3.21. Note. For a given $m = w$ new suitable measures may be constructed, if to use images of measures $m^g(E) = m(g^{-1}(E))$ such that for a diffeomorphism $g \in \text{Diff}^1(\mathbf{K})$ (see §A.3) we have $m^{g^{-1}}(dx)/m(dx) = |(g'(g^{-1}(x)))|_{\mathbf{K}}$, where $|*|_{\mathbf{K}} = \text{mod}_{\mathbf{K}}(*)$ is the modular function of the field $\mathbf{K}$ associated with the Haar measure on $\mathbf{K}$, at the same time $|*|_{\mathbf{K}}$ is the multiplicative norm in $\mathbf{K}$ consistent with its uniformity [44]. Indeed, for $\mathbf{K}$ and $X = \mathbf{K}^j$ with $j \in \mathbf{N}$ and the Haar measure $v = w$ or $v = w'$ on X , $v_X := v$ with values either in $\mathbf{R}$ or in $\mathbf{K}_s$ for $s \neq p$ and for a function $f \in L^1(X, v, \mathbf{R})$ or $f \in L(X, v, \mathbf{K}_s)$, respectively, due to Lemma 4 and Theorem 4 §I.2[44], Theorems 9.2 and A.7[35] we have: $\int_{g(A)} f(x)v(dx) = \int_A f(g(y))|g'(y)|_{\mathbf{K}}v(dy)$, where $\text{mod}_{\mathbf{K}}(\lambda)v(dx) := v(\lambda dx)$, $\lambda \in \mathbf{K}$. For a construction of new measures images of v may be used, for example, for analytic h on $\mathbf{K}$ (see 43.1 [35]) with $g = Ph$, where P is the antidifferentiation such that $g' = h$. Defining it at first locally we can then extend it on all $\mathbf{K}$.

Henceforward, quasi-invariant measure μ on $Bf(c_0(\omega_0, \mathbf{K}))$ constructed with the help of projective limits or sequences of weak distributions of probability measures $(\mu_{H(n)} : n)$ are considered, for example, as in Theorem 3.20 such that

(i) $\mu_{H(n)}(dx) = f_{H(n)}(x)v_{H(n)}(dx)$, $\dim_{\mathbf{K}} H(n) = m(n) < \aleph_0$ for each $n \in \mathbf{N}$, where $f_{H(n)} \in L^1(H(n), v_{H(n)}, \mathbf{R})$, $H(n) \subset H(n+1) \subset \dots$, $cl(\bigcup_n H(n)) = c_0(\omega_0, \mathbf{K})$, if it is not specified in another manner.

For real-valued probability quasi-invariant measures for a sequence of a weak distribution $(\mu_{H(n)})$ generated by μ in view of Proposition 11 §VII.1.9 [5], Lemma 2.3 and Definition 3.14 this condition is satisfied, since $cl(J_\mu) =$

$c_0(\omega_0, \mathbf{K})$.

As will be seen below such measures μ are quasi-invariant relative to families of the cardinality $\mathfrak{c} = \text{card}(\mathbf{R})$ of linear and non-linear transformations $U : X \rightarrow X$. Moreover, for each V open in X we have $\mu(V) > 0$, when $f_{H(n)}(x) > 0$ for each $n \in \mathbf{N}$ and $x \in H(n)$.

Let μ be a probability quasi-invariant measure satisfying (i) and $(e_j : j)$ be an orthonormal basis in M_μ , $H(n) := \text{sp}_{\mathbf{K}}(e_1, \dots, e_n)$, we denote by

$\hat{\rho}_\mu(a, x) = \hat{\rho}(a, x) = \lim_{n \rightarrow \infty} \rho^n(P_n a, P_n x)$, $\rho^n(P_n a, P_n x) := f_{H(n)}(P_n(x - a)) / f_{H(n)}(P_n x)$ for each a and x for which this limit exists and $\hat{\rho}(a, x) = 0$ in the contrary case, where $P_n : X \rightarrow H(n)$ are chosen consistent projectors. Let $\rho(a, x) = \hat{\rho}(a, x)$, if $\int \hat{\rho}(a, x) \mu(dx) = 1$, $\rho(a, x)$ is not defined when $\hat{\rho}(a, x) \neq 1$. If for some another basis $(\tilde{e}_j : j)$ and $\tilde{\rho}$ is accomplished

(ii) $\mu(S) = 1$, $S := \bigcap_{a \in M_\mu} [x : \rho(a, x) = \tilde{\rho}(a, x)]$, then $\rho(a, x)$ is called regularly dependent from a basis.

3.22. Lemma. *Let μ be a probability measure, $\mu : Bf(X) \rightarrow \mathbf{R}$, X be a Banach space over $\mathbf{K}$, suppose that for each basis $(\tilde{e}_j : j)$ in M_μ a quasi-invariance factor $\tilde{\rho}$ satisfies the following conditions:*

(1) *if $\tilde{\rho}(a_j, x)$, $j = 1, \dots, N$, are defined for a given $x \in X$ and for each $\lambda_j \in \mathbf{K}$ then a function $\tilde{\rho}(\sum_{j=1}^N \lambda_j a_j, x)$ is continuous by λ_j , $j = 1, \dots, N$;*

(2) *there exists an increasing sequence of subspaces $H(n) \subset M_\mu$, $\text{cl}(\bigcup_n H(n)) = X$, with projectors $P_n : X \rightarrow H(n)$, $B \in Bf(X)$, $\mu(B) = 0$ such that $\lim_{n \rightarrow \infty} \tilde{\rho}(P_n a, x) = \tilde{\rho}(a, x)$ for each $a \in M_\mu$ and $x \notin B$ for which is defined $\rho(a, x)$. Then $\rho(a, x)$ depends regularly from the basis.*

Proof. There exists a subset S dense in each $H(n)$, hence $\mu(B') = 0$ for $B' = \bigcup_{a \in S} [x : \rho(a, x) \neq \tilde{\rho}(a, x)]$. From (1) it follows that $\tilde{\rho}(a, x) = \rho(a, x)$ on each $H(n)$ for $x \notin B'$. From $\text{sp}_{\mathbf{K}} S \supset H(n)$ and (2) it follows that $\rho(a, x) = \tilde{\rho}(a, x)$ for each $a \in M_\mu$ and $x \in X \setminus (B' \cup B)$, consequently, Condition 3.21(ii) is satisfied.

3.23. Lemma. *If a probability quasi-invariant measure $\mu : Bf(X) \rightarrow \mathbf{R}$ satisfies Condition 3.21(i), then there exists a compact operator $T : X \rightarrow X$ such that $M_\mu \subset (TX)^\sim$, where X is the Banach space over $\mathbf{K}$.*

Proof. This follows from Theorem 3.9 (see also Part II).

3.24. Let X be a Banach space over $\mathbf{K}$, $|\ast|_{\mathbf{K}} = \text{mod}_{\mathbf{K}}(\ast)$, $U : X \rightarrow X$ be an invertible linear operator, $\mu : Bf(X) \rightarrow \mathbf{R}$ be a probability quasi-invariant measure.

The uniform convergence of a (transfinite) sequence of functions on $Af(V, \nu)$ -compact subsets of a topological space V is called the Egorov condition,

where ν is a measure on V .

Theorem. *Let pairs $(x - Ux, x)$ and $(x - U^{-1}, x)$ be in $\text{dom}(\tilde{\rho}(a, x))$, where $\text{dom}(f)$ denotes a domain of a function f , $\tilde{\rho}(x - Ux, x) > 0$, $\tilde{\rho}(x - U^{-1}x, x) > 0 \pmod{\mu}$. Then $\nu \sim \mu$ and*

$$(i) \quad \nu(dx)/\mu(dx) = |\det(U)|_{\mathbf{K}} \tilde{\rho}(x - U^{-1}x, x),$$

if ρ depends regularly from the base, then $\tilde{\rho}$ may be substituted by ρ in formula (i), where $\nu(A) := \mu(U^{-1}A)$ for each $A \in Bf(X)$.

Proof. In view of Theorem 3.9 (and Lemma 3.23) there exists a compact operator $T : X \rightarrow X$ such that $M_\mu \subset (TX)^\sim$, consequently, $(U - I)$ is a compact operator, where I is the identity operator. From the invertibility of U it follows that $(U^{-1} - I)$ is also compact, moreover, there exists $\det(U) \in \mathbf{K}$. Let g be a continuous bounded function, $g : \tilde{H}(n) \rightarrow \mathbf{R}$, whence $\int_X \phi(x) \nu(dx) = \int_{\tilde{H}(n)} g(x) [f_{H(n)}(U^{-1}x) / f_{\tilde{H}(n)}(x)] |\det(U_n)|_{\mathbf{K}} \mu_{\tilde{H}(n)}(dx)$, for $\phi(x) = g(\tilde{P}_n x)$, where subspaces exist such that $\tilde{H}(n) \subset X$, $(U^{-1} - I)\tilde{H}(n) \subset \tilde{H}(n)$, $\text{cl}(\bigcup_n \tilde{H}(n)) = X$, $U_n := \hat{r}_n(U)$, $r_n = \tilde{P}_n : X \rightarrow \tilde{H}(n)$ (see §§3.8 and 3.21), $\tilde{H}(n) \subset \tilde{H}(n+1) \subset \dots$ due to compactness of $(U - I)$. In view of the Fatou theorem $J_m \geq J_{m,\rho}$, where $J_m := \int_X g(\tilde{P}_m x) \nu(dx)$ and $J_{m,\rho} := \int_X g(\tilde{P}_m x) \tilde{\rho}(x - U^{-1}x, x) |\det(U)|_{\mathbf{K}} \mu(dx)$. Indeed, there exists n_0 such that $|u(i, j) - \delta_{i,j}| \leq 1/p$ for each i and $j > n_0$, consequently, $|\det(U_n)|_{\mathbf{K}} = |\det(U)|_{\mathbf{K}}$ for each $n > n_0$, where $\delta_{i,j}$ is the Kronecker delta-symbol. Then due to the Fubini Theorem there exists $\lim_{n \rightarrow \infty} [\mu_{\tilde{H}(n)}(d\tilde{P}_n x) / \nu_{\tilde{H}(n)}(d\tilde{P}_n x)] = \mu(dx) / \nu(dx) \pmod{\nu}$. Further analogously to the proofs of Theorems 1 and 2 §25[38].

3.25. Let X be a Banach space over $\mathbf{K}$, $|*|_{\mathbf{K}} = \text{mod}_{\mathbf{K}}(*)$ with a probability quasi-invariant measure $\mu : Bf(X) \rightarrow \mathbf{R}$, also let U fulfils the following conditions:

- (i) $U(x)$ and $U^{-1}(x) \in C^1(X, X)$ (see §A.3);
- (ii) $(U'(x) - I)$ is compact for each $x \in X$;
- (iii) $(x - U^{-1}(x))$ and $(x - U(x)) \in J_\mu$ for μ -a.e. $x \in X$;
- (iv) for μ -a.e. x pairs $(x - U(x); x)$ and $(x - U^{-1}(x); x)$

are contained in a domain of $\rho(z, x)$ such that $\rho(x - U^{-1}(x), x) \neq 0$, $\rho(x - U(x), x) \neq 0 \pmod{\mu}$;

$$(v) \quad \mu(S') = 1,$$

where $S' := ([x : \rho(z, x) \text{ is defined and continuous by } z \in L])$ for each finite-dimensional $L \subset J_\mu$;

(vi) there exists S with $\mu(S) = 0$ and for each

$x \in X \setminus S$ and for each z for which there exists $\rho(z, x)$ satisfying the following condition: $\lim_{n \rightarrow \infty} \rho(P_n z, x) = \rho(z, x)$ and the convergence is uniform for each finite-dimensional $L \subset J_\mu$ by z in $L \cap [x \in J_\mu : \|x\| \leq c]$, where $c > 0$, $P_n : X \rightarrow H(n)$ are projectors onto finite-dimensional subspaces $H(n)$ over $\mathbf{K}$ such that $H(n) \subset H(n+1)$ for each $n \in \mathbf{N}$ and $\bigcup \{H(n) : n\} = X$;

(vii) there exists n for which for all $j > n$ and $x \in X$ mappings

$V(j, x) := x + P_j(U^{-1}(x) - x)$ and $U(j, x) := x + P_j(U(x) - x)$ are invertible and $\lim_j |\det U'(j, x)| = |\det U'(x)|$, $\lim_j |\det V'(j, x)| = 1/|\det U'(x)|$.

Theorem. *The measure $\nu(A) := \mu(U^{-1}(A))$ is equivalent to μ and*

$$(i) \quad \nu(dx)/\mu(dx) = |\det U'(U^{-1}(x))|_{\mathbf{K}} \rho(x - U^{-1}(x), x).$$

Proof. I. Let at first U be linear. In general, for a linear operator U with compact $B = U - I$ there is the following decomposition $U = SCDE$, where C^t and E are upper triangular infinite matrices, $D = \text{diag}\{d(j) : j \in \mathbf{N}\}$, operators $C - I$, $D - I$ and $E - I$ are compact in the corresponding orthonormal basis $\{e_j : j\}$ in X , S transposes a finite number of vectors in orthonormal basis (see §A.2). Moreover, there are $\det(C) = \det(E) = 1$, $\det(U) = \det(D) \neq 0$.

II. Let V_n be a diagonal (or upper or lower triangular) operator on X such that $(V_n - I)(X) = P_n L$, where $\dim_{\mathbf{K}} L = k < \aleph_0$, $\lim_{n \rightarrow \infty} \|V_n - U_1\| = 0$, U_1 is a diagonal (or lower or upper triangular) operator, there exists n_0 such that $\|e_j - P_n e_j\| \leq 1/p$ for (e_j) in L , consequently, $\|\sum_j \lambda_j P_n e_j\| = \max_j |\lambda_j|$ for $\lambda_j \in \mathbf{K}$ and $\dim_{\mathbf{K}} P_n L = k$ for $n > n_0$, in addition, $\lim_{n \rightarrow \infty} \sup_{x \in L, \|x\| \leq 1} \|x - P_n x\| = 0$. Then $\lim_n \tilde{P}_n^{-1}(x - V_n^{-1}x) = x - U_1^{-1}x$, where $\tilde{P}_n x = P_n x \in P_n L$ for each $x \in L$. Due to Conditions (vi, vii) we get $\lim_n \rho(x - V_n^{-1}x, x) = \rho(x - U_1^{-1}x, x) \pmod{\mu}$.

From the Fatou theorem and §3.21 it follows that, $J_1 \geq J_{1,\rho}$, where $J_1 = \int_X f(U_1 x) \mu(dx)$, $J_{1,\rho} := \int_X f(x) \rho(x - U_1^{-1}x, x) |\det U_1|_{\mathbf{K}} \mu(dx)$ for continuous bounded function $f : X \rightarrow [0, \infty)$. Analogously for U_1^{-1} instead of U_1 . Using

instead of f the function $\Phi(U_1^{-1}x) := f(x)\rho_\mu(x - U_1^{-1}x, x)$ and Properties 3.7 we get that $\rho_\mu(U_1x - x, U_1x)\rho_\mu(x - U_1x, x) = 1 \pmod{\mu}$. Therefore, for $U = U_1U_2$ with diagonal U_1 and upper triangular U_2 and lower triangular U_3 operators with finite-dimensional over $\mathbf{K}$ subspaces $(U_j - I)X$, $j = 1, 2, 3$, the following equation is accomplished

$$\int_X f(Ux)\mu(dx) = \int_X f(x)\rho_\mu(x - U^{-1}x, x)|\det U|_{\mathbf{K}}\mu(dx).$$

If $(S^{-1}U - I)X = L$, then from the decomposition given in (I) $U = SU_2U_1U_3$, we have $(U_j - I)X = L$, $j = 1, 2, 3$ due to formulas from §A.1, since corresponding non-major minors are equal to zero.

(III). If U is an arbitrary linear operator satisfying the conditions of this theorem, then from (iv-vi) and (I, II) for each continuous bounded function $f : X \rightarrow [0, \infty)$ we have $J \geq J\rho$, where $J := \int_X f(U(x))\mu(dx)$ and $J_\rho := \int_X f(x)\rho_\mu(x - U^{-1}(x), x)|\det U|_{\mathbf{K}}\mu(dx)$. Analogously for U^{-1} , moreover, $\rho(x - U^{-1}(x), x)|\det U|_{\mathbf{K}} =: h(x) \in L^1(X, \mu, \mathbf{R})$, $h(x) > 0 \pmod{\mu}$, since there exists $\det U$.

(IV). Now let U be linear and $(U - I)(X) = L$, $\dim_{\mathbf{K}} L = k < \aleph_0$, $L \subset J_\mu$. Suppose U is polygonal, which means that there exists a partition $X = \cup\{Y(i) : i = 1, \dots, l\}$, $U(x) = a(i) + V(i)x$ for $x \in Y(i)$, where $Y(i)$ are closed subsets, $\text{Int}Y(i) \cap \text{Int}Y(j) = \emptyset$ for each $i \neq j$, $a(i) \in X$ and $V(i)$ are linear operators. Then U^{-1} is also polygonal, $U'(x) = V(j)$ for $x \in Y(j)$ and $\int_X f(a(i) + V(i)x)\mu(dx) = \int f(a(i) + x)\rho_\mu(x - V^{-1}(i)x, x) \times |\det(V(i))|_{\mathbf{K}}\mu(dx)$ for each real Borel measurable function f and each i . From $a(j) \in M_\mu$ and §3.7 we get $\int_X f(a(j) + V(j)x)\mu(dx) = \int_X f(x)\rho(x - V(j)^{-1}(x - a(j)), x)|\det V(j)|_{\mathbf{K}}\mu(dx)$. Let $H_{k,j} := [x \in X : V(k)^{-1}x = V(j)^{-1}x]$, assume without loss of generality that $V(k) \neq V(j)$ or $a(k) \neq a(j)$ for each $k \neq j$, since $Y(k) \neq Y(j)$ (otherwise they may be united). Therefore, $H_{k,j} \neq X$. If $\mu(H_{k,j}) > 0$, then from $X \ominus H_{k,j} \supset \mathbf{K}$ it follows that $M_\mu \subset H_{k,j}$, but $\text{cl}(H_{k,j}) = H_{k,j}$ and $\text{cl}(M_\mu) = X$. This contradiction means that $\mu(A) = 0$, where $A = [x : V(k)^{-1}(x - a(k)) = V(j)^{-1}(x - a(j))]$. Then $\int_X f(U(x))\mu(dx) = \int_X f(x)\rho(x - U^{-1}(x), x)|\det U'(x)|_{\mathbf{K}}^{-1}\mu(dx)$.

(V). In view of the Hahn-Banach theorem for $\mathbf{K}$ [33, 34] there are linear continuous functionals $\tilde{e}_j$ such that there exists orthonormal basis $(e_j : j)$ in X with $\tilde{e}_i(e_j) = \delta_{i,j}$. Let $s(i, j; x) := \tilde{e}_i(U(j, x) - x)$ and $s(i; x) = \tilde{e}_i(U(x) - x) = \lim_j s(i, j; x)$ (lim is taken in $C(X, \mathbf{K})$), consequently, $\det U'(j, x) = \det(ds(i, j; x)e(k) : i, k = 1, \dots, j)$. Then for the construction of the sequence

$\{U(j, *) : j\}$ it is sufficient to construct a sequence of polygonal functions $\{a(i, j; x)\}$, that is $a(i, j; x) = l_k(i, j)(x) + a_k$ for $x \in Y(k)$, where $l_k(i, j)$ are linear functionals, $a_k \in \mathbf{K}$, $Y(k)$ are closed in X , $Int(Y(j)) \cap Int(Y(k)) = \emptyset$ for each $k \neq j$, $\bigcup_{k=1}^m Y(k) = X$, $m < \aleph_0$. For each $c > 0$ there exists $V_c \subset X$ with $\mu(X \setminus V_c) < c$, the functions $s(i, j; x)$ and $(\bar{\Phi}^1 s(i, j; *))(x, e(k), t)$ are equiuniformly continuous (by $x \in V_c$ and by $i, j, k \in \mathbf{N}$) on V_c . Choosing $c = c(n) = 1/n$ and using δ -nets in V_c we get a sequence of polygonal mappings $(W_n : n)$ converging by its matrix elements in $L^1(X, \mu, \mathbf{R})$, from Condition (i) follows that it may be chosen equicontinuous for matrix elements $s(i, j; x)$, $ds(i, j; x)$ and $s(i, P_j x)$ by i, j (the same is true for U^{-1}).

Indeed, for V_c there is $\delta > 0$ such that $|s(i, j; x') - s(i, j; x)| < c$ and $|(\bar{\Phi}^1 s(i, j; *))(x, e(k), t) - (\bar{\Phi}^1 s(i, j; *))(x', e(k), t')| < c$ (see Theorem 2.11.I [22]) for each $x, x' \in V_c$ with $|x - x'| < \delta$, $|t - t'| \leq 1$ and $i, j, k \in \mathbf{N}$. Then we can choose in V_c a finite δ -net $x_1, \dots, x_r$ and define $l_q(i, j; x) = s(i, j; x_q) + (ds(i, j; x_q))(x - x_q)$ and apply the non-Archimedean variant of the Taylor theorem (see §29.4[35] and Theorem A.5).

Then calculating integrals as above for W_n with functions f , using the Fatou theorem we get the inequalities analogous to written in (III) for J and J_ρ of the general form. From $\nu(dx)/\mu(dx) > 0 \pmod{\mu}$ we get the statement of the theorem (see also Theorem 1 §26[38] in the classical case), since

$$\int_X f(A(j, x))\mu(dx) = \int_X f(x)\rho_\mu(P_j(x - A(x)), x)|\det(A(j, x'))|_{\mathbf{K}}^{-1}\mu(dx),$$

where $A = U$, $A(j, x) = U(j, x)$ or $A = U^{-1}$, $A(j, x) = V(j, x)$.

3.26. Note. For a linear invertible operator U Condition 3.25(i) is satisfied automatically, (ii) follows from (iii), hence from (iii) also follows (vii) (see the proof of Theorem 3.24).

3.27. Examples. Let X be a Banach space over the field $\mathbf{K}$ with the valuation group $\Gamma_{\mathbf{K}} = \Gamma_{\mathbf{Q}_p}$. We consider a diagonal compact operator $T = \text{diag}(t_j : j \in \mathbf{N})$ in a fixed orthonormal basis (e_j) in X such that $\ker T := T^{-1}0 = \{0\}$. Let $\nu_j(dx_j) = C(\xi_j)\exp(-|(x_j - x_j^0)/\xi_j|_p^q)v(dx_j)$ for the Haar measure $v : Bf(\mathbf{K}) \rightarrow [0, \infty)$. We choose constant functions $C(\xi_j)$ and $C'(\xi_j)$ such that ν_j be a probability measure, where $x^0 = (x_j^0 : j) \in X$, $x = (x_j : j) \in X$, $x_j \in \mathbf{K}$. Particularly, for $q = 2$: $\int_{\mathbf{K}}[(\chi_e(x_j - x_j^0) - \chi_e(-x_j + x_j^0))/(2i)]\nu_j(dx_j) = 0$, since ν_j is symmetric and $[\chi_e(x) - \chi_e(-x)]$ is the odd function, where χ_e is the same character as in §2.6. Then $\int_{\mathbf{K}} |x_j - x_j^0|^2 \nu_j(dx_j) = |\xi_j|^2$ due to the formula of changing variables in §3.21

and differentiation by the real parameter b of the following integral $I(b) = \int_{\mathbf{K}} \exp(-|x|_p^2 b^2) v(dx)$, then we take $1/|\xi| = b \in \Gamma_{\mathbf{K}}$. Therefore, x_j^0 and $|\xi_j|^2$ have in some respect a meaning looking like the mean value and the variance.

With the help of products $\bigotimes_{j=1}^{\infty} \nu_j(dx_j)$ as in §3.20 we can construct a probability quasi-invariant measure μ^T on X with values in $[0, 1]$ or $\mathbf{C}_s$, since $cl(TX)$ is compact in X and $sp_{\mathbf{K}}(e_j : j) =: H \subset J_{\mu}$. From $\bigcap_{\lambda \in B(\mathbf{K}, 0, 1) \setminus 0} cl(\lambda TX) = \{0\}$ we may infer that for each $c > 0$ there exists a compact $V_c(\lambda) \subset X$ such that $\mu(X \setminus V_c(\lambda)) < c$ and $\bigcap_{\lambda \neq 0} V_c(\lambda) = \{0\}$, consequently, $\lim_{|\lambda| \rightarrow 0} \int_X f(x) \mu^{\lambda T}(dx) = f(0) = \delta_0(f)$, hence $\mu^{\lambda T}$ is weakly converging to δ_0 whilst $|\lambda| \rightarrow 0$ for the space of bounded continuous functions $f : X \rightarrow \mathbf{R}$.

From Theorem 3.4 we conclude that from $\sum_{j=1}^{\infty} |y_j/\xi_j|_p^q < \infty$ it follows $y \in J_{\mu^T}$. Then for a linear transformation $U : X \rightarrow X$ from $\sum_j |\tilde{e}_j(x - U(x))/\xi_j|_p^2 < \infty$ it follows that $x - U(x) \in J_{\mu}$ and a pair $(x - U(x), x) \in dom(\rho(a, z))$. Moreover, for ρ corresponding to μ^T conditions (v) and (vi) in §3.25 are satisfied. Therefore, for such y and $S \in Af(X, \mu)$ a quantity $|\mu(ty + S) - \mu(S)|$ is of order of smallness $|t|^q$ whilst $t \rightarrow 0$, hence they are pseudo-differentiable of order b for $0 < Re(b) < 2$ (see also §4 below).

Analogues of real-valued Wiener measures on spaces of functions from a non-Archimedean Banach space into $\mathbf{C}$ are given in [2], that are quasi-invariant under some suitable choice.

4 Pseudo-differentiable measures.

4.1. Definition and notes. A function $f : \mathbf{K} \rightarrow \mathbf{R}$ is called pseudo-differentiable of order b , if there exists the following integral: $PD(b, f(x)) := \int_{\mathbf{K}} [(f(x) - f(y)) \times g(x, y, b)] dv(y)$. We introduce the following notation $PD_c(b, f(x))$ for such integral by $B(\mathbf{K}, 0, 1)$ instead of the entire $\mathbf{K}$. Where $g(x, y, b) := |x - y|^{-1-b}$ with the corresponding Haar measure v with values in $\mathbf{R}$, where $b \in \mathbf{C}$ and $|x|_{\mathbf{K}} = p^{-ord_p(x)}$.

Obviously, the definitions of differentiability of measures can not be transferred from [4, 8] onto the case considered here. This is the reason why the notion of pseudo-differentiability is introduced here. A quasi-invariant measure μ on X is called pseudo-differentiable for $b \in \mathbf{C}$, if there exists $PD(b, g(x))$ for $g(x) := \mu(-xz + S)$ for each $S \in Af(X, \mu)$ with $|m|(S) < \infty$ and each $z \in J_{\mu}^b$, where J_{μ}^b is a $\mathbf{K}$ -linear subspace dense in X . For a fixed $z \in X$ such measure is called pseudo-differentiable along z .

For a one-parameter subfamily of operators $B(\mathbf{K}, 0, 1) \ni t \rightarrow U_t : X \rightarrow X$ quasi-invariant measure μ is called pseudo-differentiable for $b \in \mathbf{C}$, if for each S the same as above there exists $PD_c(b, g(t))$ for a function $g(t) := \mu(U_t^{-1}(S))$, where X may be also a topological group G with a measure quasi-invariant relative to a dense subgroup G' (see [22, 23, 26]).

4.2. Let μ , X , and ρ be the same as in Theorem 3.24 and F be a non-Archimedean Fourier transform defined in [43, 34].

Theorems. (1) $g(t) := \rho(z+tw, x)j(t) \in L^1(v, K \rightarrow \mathbf{C}) := V$, moreover, there exists $F(g) \in C^0(\mathbf{K}, \mathbf{C})$ and $\lim_{|t| \rightarrow \infty} F(g)(t) = 0$ for μ and v with values in $\mathbf{R}$, where z and $w \in J_\mu$, $t \in \mathbf{K}$, $j(t)$ is the characteristic function of a compact subset $W \subset \mathbf{K}$. In general, may be $k(t) := \rho(z+tw, x) \notin V$.

(2) Let $g(t) = \rho(z+tw, x)j(t)$ with clopen subsets W in $\mathbf{K}$. Then there are μ , for which there exists $PD(b, g(t))$ for each $b \in \mathbf{C}$. If $g(t) = \rho(z+tw, x)$, then there are probability measures μ , for which there exists $PD(b, g(t))$ for each b with $0 < \text{Re}(b)$ or $b = 0$.

(3) Let $S \in Af(X, y)$, $|\mu|(S) < \infty$, then for each $b \in U := \{b' : \text{Re } b' > 0 \text{ or } b' = 0\}$ there is a pseudo-differentiable quasi-invariant measure μ .

Proof. We consider the following additive compact subgroup $G_T := \{x \in X \mid \|x(j)\| \leq p^{k(j)} \text{ for each } j \in \mathbf{N}\}$ in X , where $T = \text{diag}\{d(j) \in K : |d(j)| = p^{-k(j)} \text{ for each } j \in \mathbf{N}\}$ is a compact diagonal operator. Then μ from Theorem 3.20 is quasi-invariant relative to the following additive subgroup $S_T := G_T + H$, where $H := sp_{\mathbf{K}}\{e(j) : j \in \mathbf{N}\}$. Moreover, for each z and $w \in G_T$ and $R > 0$ there is $c > 0$ such that $\rho(z+uw, x) = \rho(z+sw, x)$ for $|u-s| < c$ and $x \in B(X, 0, R)$, if all functions f_j in the proof of Theorem 3.20 are locally constant (that is, f are defined on $\mathbf{K}e_j \subset X$). In general, for each $b \in \mathbf{C}$ we can choose a sequence $h_j(x)$ with $\sum_{j=0}^{\infty} \sup_{x \in \mathbf{K}} (|h_j(x)/(h_j(x) + f_j(x))| r_j(x)) < c'$ for suitable fixed $c' > 0$ and functions $r_j : \mathbf{K} \rightarrow [0, \infty)$ with $\lim_{|x| \rightarrow 0} r_j(x) = \infty$. Carrying out calculations and using the fact that μ is the probability quasi-invariant measure we get the pseudo-differentiability of μ . Using the Riemann-Lebesgue theorem from [43] we get the statement of this theorem for $F(g)$.

4.3. Let X be a Banach space over $\mathbf{K}$, $b_0 \in \mathbf{R} \cup \{+\infty\}$ and suppose that the following conditions are satisfied:

- (1) $T : X \rightarrow X$ is a compact operator with $\ker(T) = \{0\}$;
- (2) a mapping $\tilde{F}$ from $B(\mathbf{K}, 0, 1)$ to $C_T(X) := \{U : U \in C^1(X, X) \text{ and } (U'(x) - I) \text{ is a compact operator for each } x \in X, \text{ there is } U^{-1} \text{ satisfying the same conditions as } U\}$ is given;

(3) $\tilde{F}(t) = U_t(x)$ and $\bar{\Phi}^1 U_t(x + h, x)$ are continuous by t , that is, $\tilde{F} \in C^1(B(\mathbf{K}, 0, 1), C_T(X))$;

(4) there is $c > 0$ such that $\|U_t(x) - U_s(x)\| \leq \|Tx\|$ for each $x \in X$ and $|t - s| < c$;

(5) for each $R > 0$ there is a finite-dimensional over $\mathbf{K}$ subspace $H \subset X$ and $c' > 0$ such that $\|U_t(x) - U_s(x)\| \leq \|Tx\|/R$ for each $x \in X \ominus H$ and $|t - s| < c'$ with (3 – 5) satisfying also for U_t^{-1} .

Theorem. *On X there are probability quasi-invariant measures μ which are pseudo-differentiable for each $b \in \mathbf{C}$ with $\mathbf{R} \ni \operatorname{Re}(b) \leq b_0$ relative to a family U_t .*

Proof. From Conditions (2,3) it follows that there is $c > 0$ such that $|\det(U'_t(x))| = |\det(U'_s(x))|$ for μ -a.e. $x \in X$ and all $|t - s| < c$, where quasi-invariant and pseudo-differentiable measures μ on X relative to S_T may be constructed as in the proof of Theorems 3.20 and 4.2. From (1 – 5) it follows that conditions of Theorem 3.25 are satisfied for each U_t . From (3, 5) it follows that for each $R > 0$ and $b > 0$ there exists $c' > 0$ such that $|\rho_\mu(x - U_t^{-1}(x), x) - \rho_\mu(x - U_t^{-1}(x), x)| \leq b$ for each $|t - s| < c'$ and $x \in B(X, 0, R)$. If during construction of μ to use only locally constant $f_j(x_j)$ with $h_j = 0$, then we can take $\infty \geq b_0 \geq 0$. Indeed, let $T = SCDE$ be a decomposition from the appendix and $\mu^A(\tilde{S}) = \mu(E\tilde{S})$ with $A = E^{-1}$ (or C), $\tilde{S} \in Bf(X)$, then $\rho_{\mu^A}(x - U_t^{-1}(x), x) = \rho_\mu(Ex - U_t^{-1}(Ex), Ex)$ and $c(x) := \|U_t^{-1}(Ex) - U_s^{-1}(Ex)\| \leq \|TEEx\|$ for $x \in X$ and $c(x) \leq \|TEEx\|/R$ for $x \in X \ominus H$ (for each $R > 0$ there is $H \subset X$ with $\dim_{\mathbf{K}} H < \infty$), due to compactness of $E - I$ and $C - I$ we can choose H such that $A^{-1}H = H$, $AH = H$. We use also images of quasi-invariant measure constructed with the help of compact diagonal operator $(D' - I)$, where $|D_j - 1| < |D'_j - 1|p^{k(j)}$, $\lim_j k(j) = -\infty$. Then $\sup\{\|U_t(x) - U_t(0)\| : x \in B(X, 0, R)\} \leq \|\bar{\Phi}^1 U_t\|_{B(X, 0, R)} \times \|x\|$. From the fact that H is finite-dimensional and the existence of the orthogonal projector (that is, corresponding to the decomposition into the direct sum) $\pi_H : X \rightarrow H$ it follows that $\|U_t^a(e_j) - U_s^a(e_j)\| \leq b_j$ for each $j \in \mathbf{N}$ and each $|t - s| < c'$, where $\lim_{j \rightarrow \infty} b_j = 0$, $a = 1$ or $a = -1$. This guarantees pseudo-differentiability of μ .

4.4. Let X be a Banach space over $\mathbf{K}$, μ be a probability quasi-invariant measure $\mu : Bf(X) \rightarrow \mathbf{R}$, that is pseudo-differentiable for a given b with $\operatorname{Re}(b) > 0$, $C_b(X)$ be a space of continuous bounded functions $f : X \rightarrow \mathbf{R}$ with $\|f\| := \sup_{x \in X} |f(x)|$.

Theorem. *For each $a \in J_\mu$ and $f \in C_b(X)$ is defined the following*

integral:

$$(i) \quad l(f) = \int_{\mathbf{K}} \left[\int_X f(x) [\mu(-\lambda a + dx) - \mu(dx)] g(\lambda, 0, b) v(d\lambda) \right]$$

and there exists a measure $\nu : Bf(X) \rightarrow \mathbf{C}$ with a bounded variation (for $b \in \mathbf{R}$ this ν is a mapping from $Bf(X)$ into $\mathbf{R}$) such that

$$(ii) \quad l(f) = \int_X f(x) \nu(dx),$$

where v is the Haar measure on $\mathbf{K}$ with values in $[0, \infty)$, moreover, ν is independent from f and may be dependent on $a \in J_\mu$. We denote $\nu =: \tilde{D}_a^b \mu$.

Proof. From Definition 4.1 and the Fubini and Lebesgue theorems it follows that there exists

$$\lim_{j \rightarrow \infty} \int_{\mathbf{K} \setminus B(\mathbf{K}, 0, p^{-j})} \left[\int_X (f(x + \lambda a) - f(x)) g(\lambda, 0, b) \mu(dx) \right] v(d\lambda) = l(f),$$

that is (i) exists. Let

$$(iii) \quad l_j(V, f) := \int_{\mathbf{K} \setminus B(\mathbf{K}, 0, p^{-j})} \left[\int_V f(x) (\mu(-\lambda a + dx) - \mu(dx)) g(\lambda, 0, b) v(d\lambda) \right],$$

where $V \in Bf(X)$. The proofs of §§2 and 3 in Theorem 1 §21[38] may be slightly transformed for the considered here case of the Banach space X over $\mathbf{K}$ instead of a Hilbert space over $\mathbf{R}$. Hence for each $c > 0$ there exists a compact $V_c \subset X$ with $|\nu_\lambda|(X \setminus V_c) < c$ for each $|\lambda| > 0$, where $\nu_\lambda(A) := \int_{\mathbf{K} \setminus B(\mathbf{K}, 0, |\lambda|)} [\mu(-\lambda' a + A) - \mu(A)] g(\lambda', 0, b) v(d\lambda')$ for $A \in Bf(X)$. Also there are $\delta > 0$ and V_c such that $-\lambda' a + V_c \subset V_c$ and $\| [X \setminus V_c] \triangle (-\lambda' a + (X \setminus V_c)) \|_\mu = 0$ for each $|\lambda'| < \delta$ (see also Theorem 7.22 [34]), where $A \triangle B := (A \setminus B) \cup (B \setminus A)$.

Each continuous linear functional on $C_b(V)$ for compact V has the form (ii) (see [5], A.5 [35]), that is, for $l_j(V, f)$ there exists a measure $\nu_\lambda(c, *)$. In view of the Alaoglu-Bourbaki theorem each bounded subset W in $[C_b(V_c)]'$ is precompact (i.e., $cl(W)$ is compact) in the weak topology, where $C_b(V_c)$ is separable. For each $c > 0$ we have

$$l(f) = \int_{V_c} f(x) \nu(c, dx) + O(c \times \sup_{x \notin V_c} |f(x)|)$$

such that $\nu(c, *)$ are measures on $Bf(V_c)$. Choosing $V_c \subset V_{c'}$ for $c > c'$ and defining $\nu(A) := \lim_{c \rightarrow +0} \nu(c, A \cap V_c)$ for each $A \in Bf(X)$ in view of the Radon-Nikodym theorem about convergence of measures (see [7]) we get (i) with the help of the theorem about extension of measures, since the minimal σ -field $\sigma Bco(X)$ generated by $Bco(X)$ coincides with $Bf(X)$.

4.5. Theorem. *If μ is probability real-valued quasi-invariant measure on a Banach space X over $\mathbf{K} = \mathbf{Q}_p$, $a \in J_\mu$ and μ is pseudo-differentiable of order 1, then for each $f \in L^\infty(X, \mu, \mathbf{R})$ with $\text{supp}(f) \subset B(X, 0, R)$, $\infty > R = R(f) > 0$ the following equality is accomplished:*

$$(i) \int_X [D_\lambda^{-1} f(x + \lambda a)]|_{\lambda=0}^{\lambda=\beta} (\tilde{D}_a^1 \mu)(dx) = \int_X [f(x + \beta a) - f(x)] \mu(dx),$$

moreover, pseudo-differential $\tilde{D}_a^1 \mu := \nu$ in the direction a from Theorem 4.4 given by formula (ii) characterizes μ uniquely.

Proof. From the restrictions on f it follows that $f(x + \lambda a) =: \psi(\lambda) \in \mathbf{E}'$, where $\mathbf{E}'$ is the topologically adjoint space to the space $\mathbf{E}$ of basic locally constant functions [43]. By formula (1.5) §9[43] for them there is the following equality $D^a D^b \psi = D^b D^a \psi = D^{a+b} \psi$ for $a + b \neq -1$, $a, b \in \mathbf{R}$ (D_λ^a denotes a pseudo-differentiation of functions by λ). In view of Theorem in §6.3[43] $\mathbf{E}' = [\phi \in \mathbf{D}' : \text{supp}(\phi) \text{ is compact}]$, where $\mathbf{D} = [\phi \in \mathbf{E} : \text{supp}(\phi) \text{ is compact}]$. Since $[\xi]_0^\beta := \xi(\beta) - \xi(0)$ for $\xi(\lambda) = D_\lambda^{-1} \psi(\lambda)$ and $(D^{-1} D \psi)(\lambda) = \psi(\lambda)$, hence due to the definition of D_λ^{-1} we have:

$$p^2(p+1)^{-1} \int_X [D_\lambda^{-1} f(x + \lambda a)]|_0^\beta (\tilde{D}_a^1 \mu)(dx) = \lim_{\alpha \rightarrow -1} \int_{\mathbf{Q}_p} \int_{\mathbf{Q}_p} (1 - p^\alpha)$$

$$(1 - p^{-1-\alpha})^{-1} p^2(1+p)^{-1} \left[\int_X (\mu(-a(\lambda' + \lambda'') + dx) - \mu(-a\lambda' + dx)) \right.$$

$$\left. |\lambda''|_p^{-2} |\lambda - \lambda'|_p^{-1-\alpha} \mu(dx) \right] v(d\lambda') v(d\lambda'').$$

In view of the Fubini and Lebesgue theorems we establish Formula (i). Taking all different $f \in L^\infty(X, \mu, \mathbf{R})$ such that $\psi(\lambda) = f(x + a\lambda)$ are locally polynomial by λ and using the non-Archimedean variant of the Stone-Weierstrass theorem by Kaplanski [35] we get that μ is characterized by the left side of (i) uniquely.

4.6. (Properties of $\tilde{D}_a^1 \mu$ for real-valued μ , see §§4.4. and 4.5, where $\mathbf{K} = \mathbf{Q}_p$.)

Proposition.1. *If μ is pseudo-differentiable in a direction $a \in J_\mu^1$, $a \neq 0$, then for each $d \in J_\mu^1$ a measure $\mu^d(A) := \mu(A - d)$, $A \in Bf(X)$, is also pseudo-differentiable in a direction a , moreover, $\tilde{D}_a^1(\mu^d) = (\tilde{D}_a^1\mu)_d$. If $a \in J_\mu^b$, then $\tilde{D}_{\lambda a}^b\mu$ exists for each $\lambda \in \mathbf{K}$.*

Proof. From §§4.4 and 4.5 it follows that

$$\begin{aligned} \int_X f(x)(\tilde{D}_a^{-1}\mu)^d(dx) &= \int_{\mathbf{K}} \int_X [f(x + \lambda a) - f(x)] |\lambda|_{\mathbf{K}}^{-2} \mu^d(dx) v(d\lambda) \\ &= \int_X f(x)(\tilde{D}_a^1(\mu^d))(dx). \end{aligned}$$

The last statement follows from Definition 4.1 with the help of §3.21.

Proposition.2. *If $a \in J_\mu^1$, $\nu = \tilde{D}_a^1\mu$ and $D_\lambda^{-1}\rho_\nu(\lambda a, x) =: \phi(x) \in L^1(X, \nu, \mathbf{R})$ by x , then $\mu^{\lambda a} \ll \nu$.*

Proof. In view of Formula 4.5.(i) the following equality is accomplished:

$$\int_X f(x)\mu^{\lambda a}(dx) = \int_X f(x)\mu(dx) + \int_X f(x)[D_\xi^{-1}\rho_\nu(\lambda a, x)]|_{\xi=0}^{\xi=\lambda} \nu(dx).$$

Taking $f(x) = \chi_A(x)$, where χ_A is a characteristic function of $A \in Bf(X)$ we get $\mu^{\lambda a}(A) = \mu(A)$ for every $|\lambda| > 0$. Therefore, considering $|\lambda| \rightarrow \infty$ we get $\mu(A) = 0$, that is, $\mu \ll \nu$ and $\mu^{\lambda a} \ll \nu$.

Proposition.3. *If a non-negative measure μ is pseudo-differentiable in a direction $a \in J_\mu^1$ and $(\tilde{D}_a^1\mu) \ll \mu$, then $l_\mu(a, x) := (\tilde{D}_a^1\mu)(dx)/\mu(dx)$ is called the logarithmic pseudo-derivative of quasi-invariant measure μ in a direction a . Then due to Formula 4.5.(i) there is the equality:*

$$\int_X [f(x + \lambda a) - f(x)]\mu(dx) = \int_X [\tilde{D}_\xi^{-1}f(x + \xi a)]|_{\xi=0}^{\xi=\lambda} l_\mu(a, x)\mu(dx).$$

We may slightly transform for the non-Archimedean case also Theorem 1 §22[38] with the help of the results of §§4.2-4.4 above.

4.7. Theorem. *Let X be a Banach space over $\mathbf{K}$, $|\cdot| = \text{mod}_{\mathbf{K}}(\cdot)$ with a probability quasi-invariant measure $\mu : Bf(X) \rightarrow \mathbf{R}$ and it is satisfied Condition 3.21(i), suppose μ is pseudo-differentiable and*

(viii) $J_b\mu \subset T''J_\mu$, $(U_t : t \in B(\mathbf{K}, 0, 1))$ is a one-parameter family of operators such that Conditions 3.25(i – vii) are satisfied with the substitution of J_μ onto J_μ^b uniformly by $t \in B(\mathbf{K}, 0, 1)$, $J_\mu \supset T'X$, where $T', T'' : X \rightarrow X$ are compact operators, $\ker(T') = \ker(T'') = 0$. Moreover, suppose that there are sequences

(ix) $[k(i, j)]$ and $[k'(i, j)]$ with $i, j \in \mathbf{N}$, $\lim_{i+j \rightarrow \infty} k(i, j) = \lim_{i+j \rightarrow \infty} k'(i, j) = -\infty$ and $n \in \mathbf{N}$ such that $|T''_{i,j} - \delta_{i,j}| < |T'_{i,j} - \delta_{i,j}| p^{k(i,j)}$, $|U_{i,j} - \delta_{i,j}| < |T''_{i,j} - \delta_{i,j}| p^{k'(i,j)}$ and $|(U^{-1})_{i,j} - \delta_{i,j}| < |T''_{i,j} - \delta_{i,j}| p^{k'(i,j)}$ for each $i + j > n$, where $U_{i,j} = \tilde{e}_i U(e_j)$, $(e_j : j)$ is orthonormal basis in X . Then for each $f \in C_b(X)$ there is defined

$$(i) \quad l(f) = \int_{B(K,0,1)} \left[\int_X f(x) [\mu(U_t^{-1}(dx)) - \mu(dx)] g(t, 0, b) v(dt) \right]$$

and there exists a measure $\nu : Bf(X) \rightarrow \mathbf{C}$ with a bounded total variation [particularly, for $b \in \mathbf{R}$ it is such that $\nu : Bf(X) \rightarrow \mathbf{R}$] and

$$(ii) \quad l(f) = \int_X f(x) \nu(dx),$$

where ν is independent from f and may be dependent on $(U_t : t)$, $\nu =: \tilde{D}_{U_*}^b \mu$.

Proof. From the proof of Theorem 3.25 it follows that there exists a sequence $U_t^{(q)}$ of polygonal operators converging uniformly by $t \in B(\mathbf{K}, 0, 1)$ to U_t and equicontinuously by indices of matrix elements in L^1 . Then there exists $\lim_{q \rightarrow \infty} \lim_{j \rightarrow \infty} \int_{B(\mathbf{K},0,1) \setminus B(\mathbf{K},0,p^{-j})} [\int_X f(U_t^{-1}(x)) - f(x)] g(t, 0, b) \mu(dx) v(dt)$ for each $f \in C_b(X)$. From conditions (viii, ix), the Fubini and Lebesgue theorems it follows that for $\nu_\lambda := \int_{B(\mathbf{K},0,1) \setminus B(\mathbf{K},0,|\lambda|)} [\mu(U_t^{-1}(A)) - \mu(A)] g(t, 0, b) v(dt)$ for $A \in Bf(X)$ for each $c > 0$ there exists a compact $V_c \subset X$ and $\delta > 0$ such that $|\nu_\lambda|(X \setminus V_c) < c$. Indeed, V_c and $\delta > 0$ may be chosen due to pseudo-differentiability of μ , §§2.35, 3.9, 3.23, Formula (i), 3.21(i) and due to continuity and boundedness (on $B(\mathbf{K}, 0, 1) \ni t$) of $|\det U'_t(U_t^{-1}(x))|_{\mathbf{K}}$ satisfying the following conditions $U_t^{-1}(V_c) \subset V_c$ and $\|(X \setminus V_c) \triangle (U_t^{-1}(X \setminus V_c))\|_\mu = 0$ for each $|t| < \delta$, since $V_c = Y(j) \cap V_c$ are compact for every j . At the same time closed subsets $Y(j) \subset X$ for $U_t^{(q)}$ may be chosen independent from t and in §2.35 operator S_c may be chosen symmetric, $TX \supset J_\mu \supset T'X$, where T is a compact operator. Evidently, conditions of type (ix) are carried out for $V(j, x)$ and $U(j, x)$ uniformly by j . Repeating proofs 3.25 and 4.4 with the use of Lemma 2.5 for the family $(U_t : t)$ we get formulas (i, ii).

5 Convergence of quasi-invariant and pseudo-differentiable measures.

Different types of convergence of measures were considered in [4, 7, 39, 40, 41]. The definitions and theorems given below are modifications of the corre-

sponding chapters of [7], but additionally taking into account the properties of quasi-invariance and pseudo-differentiability.

5.1. Definitions, notes and notations. Let S be a normal topological group with the small inductive dimension $ind(S) = 0$, S' be a dense subgroup, suppose their topologies are τ and τ' correspondingly, $\tau' \supset \tau|_{S'}$. Let G be an additive Hausdorff left- R -module, where R is a topological ring, $\mathbf{R} \supset Bf(S)$ be a σ -ring for real-valued measures, $\mathbf{M}(R, G)$ be a family of measures with values in G , $\mathbf{L}(R, G, R)$ be a family of quasi-invariant measure $\mu : \mathbf{R} \rightarrow G$ with $\rho_\mu(g, x) \times \mu(dx) := \mu^{g^{-1}}(dx) =: \mu(gdx)$, $R \times G \rightarrow G$ be a continuous left action of R on G such that $\rho_\mu(gh, x) = \rho_\mu(g, hx)\rho_\mu(h, x)$ for each $g, h \in S'$ and $x \in S$. Particularly, $1 = \rho_\mu(g, g^{-1}x)\rho_\mu(g^{-1}, x)$, that is, $\rho_\mu(g, x) \in R_o$, where R_o is a multiplicative subgroup of R . Moreover, $zy \in \mathbf{L}$ for $z \in R_o$ with $\rho_{z\mu}(g, x) = z\rho_\mu(g, x)z^{-1}$ and $z \neq 0$. We suppose that topological characters and weights S and S' are countable and each open W in S' is precompact in S . Let $\mathbf{P}''$ be a family of pseudo-metrics in G generating the initial uniformity such that for each $c > 0$ and $d \in \mathbf{P}''$ and $\{U_n \in \mathbf{R} : n \in \mathbf{N}\}$ with $\cap \{U_n : n \in \mathbf{N}\} = \{x\}$ there is $m \in \mathbf{N}$ such that $d(\mu^g(U_n), \rho_\mu(g, x)\mu(U_n)) < cd(\mu(U_n), 0)$ for each $n > m$, in addition, a limit ρ is independent μ -a.e. on the choice of $\{U_n : n\}$ for each $x \in S$ and $g \in S'$. Consider a subring $R' \subset \mathbf{R}$, $R' \supset Bf(S)$ such that $\cup \{A_n : n = 1, \dots, N\} \in R'$ for $A_n \in R'$ with $N \in \mathbf{N}$ and $S'R' = R'$. Then $\mathbf{L}(R, G, R; R') := \{(\mu, \rho_\mu(*, *)) \in \mathbf{L}(R, G, R) : \mu|_{R'} - \text{is regular and for each } s \in S \text{ there are } A_n \in R', n \in \mathbf{N} \text{ with } s = \cap (A_n : n), \{s\} \in R'\}$.

For pseudo-differentiable measures μ let $S'' \subset S'$, S'' be a dense subgroup in S , $\tau'|_{S''}$ is not stronger than τ'' on S'' and there exists a neighbourhood $\tau'' \ni W'' \ni e$ in which are dense elements lying on one-parameter subgroups $(U_t : t \in B(\mathbf{K}, 0, 1))$. We suppose that μ is induced from the Banach space X over $\mathbf{K}$ due to a local homeomorphism of neighbourhoods of e in S and 0 in X as for the case of groups of diffeomorphisms [20] such that is accomplished Theorem 4.7 for each $U_* \subset S''$ inducing the corresponding transformations on X . In the following case $S = X$ we consider $S' = J_\mu$ and $S'' = J_\mu^b$ with $Re(b) > 0$ such that $M_\mu \supset J_\mu \subset (T_\mu X)^\sim$, $J_\mu^b \subset (T_\mu^{(b)} X)^\sim$ with compact operators T_μ and $T_\mu^{(b)}$, $\ker(T_\mu) = \ker(T_\mu^{(b)}) = 0$ and norms induced by the Minkowski functional P_E for $E = T_\mu B(X, 0, 1)$ and $E = T_\mu^{(b)} B(X, 0, 1)$ respectively. We suppose further that for pseudo-differentiable measures G is equal to $\mathbf{C}$ or $\mathbf{R}$. We denote $P(R, G, R, U_*; R') := [(\mu, \rho_\mu, \eta_\mu) : (\mu, \rho_\mu) \in$

$L(\mathbf{R}, G, R; R'), \mu$ is pseudo-differentiable and $\eta_\mu(t, U_*, A) \in L^1(\mathbf{K}, v, \mathbf{C})$], where

$$\eta_\mu(t, U_*, A) = j(t)g(t, 0, b)[\mu^h(U_t^{-1}(A) - \mu^h(A))],$$

$j(t) = 1$ for each $t \in \mathbf{K}$ for $S = X$; $j(t) = 1$ for $t \in B(\mathbf{K}, 0, 1)$, $j(t) = 0$ for $|t|_{\mathbf{K}} > 1$ for a topological group S that is not a Banach space X over $\mathbf{K}$, v is the Haar measure on $\mathbf{K}$ with values in $[0, \infty)$, $(U_t : t \in B(\mathbf{K}, 0, 1))$ is an arbitrary one-parameter subgroup. On these spaces $\mathbf{L}$ (or P) the additional conditions are imposed:

(a) for each neighbourhood (implying that it is open) $U \ni 0 \in G$ there exists a neighbourhood $S \supset V \ni e$ and a compact subset V_U , $e \in V_U \subset V$, with $\mu(B) \in U$ (or in addition $\tilde{D}_{U_*}^b \mu(B) \in U$) for each B , $\mathbf{R} \ni B \in Bf(S \setminus V_U)$;

(b) for a given U and a neighbourhood $R \supset D \ni 0$ there exists a neighbourhood W , $S' \supset W \ni e$, (pseudo)metric $d \in P''$ and $c > 0$ such that $\rho_\mu(g, x) - \rho_\mu(h, x') \in D$ (or $\tilde{D}_{U_*}^b(\mu^g - \mu^h)(A) \in U$ for $A \in Bf(V_U)$ in addition for P) whilst $g, h \in W$, $x, x' \in V_U$, $d(x, x') < c$, where (a,b) is satisfied for all $(\mu, \rho_\mu) \in \mathbf{L}$ (or $(\mu, \rho_\mu, \eta_\mu) \in P$) equicontinuously in (a) on $V \ni U_t, U_t^{-1}$ and in (b) on W and on each V_U for $\rho_\mu(g, x) - \rho_\mu(h, x')$ and $\tilde{D}_{U_*}^b(\mu^g - \mu^h)(A)$.

These conditions are justified, since for a Gaussian measure ν on a Hilbert space Z over $\mathbf{C}$ or $\mathbf{R}$ (or measures given above for a non-Archimedean Banach space Z over $\mathbf{K}$) there exists a subspace Z' dense in Z such that ν is quasi-invariant relative to Z' . Moreover, in view of Theorem 26.2[38] (or Theorems 3.20, 3.25, 4.3 and 4.7 in the non-Archimedean case respectively) there exists a subspace Z'' dense in Z' such that for each $\epsilon > 0$ and each $\infty > R > 0$ there are $r > 0$ and $\delta > 0$ with $|\rho_\nu(g, x) - \rho_\nu(h, y)| < \epsilon$ for each $\|g - h\|_{Z''} + \|x - y\|_Z < \delta$, $g, h \in B(Z'', 0, r)$, $x, y \in B(Z, 0, R)$, where Z'' is the Banach space over $\mathbf{C}$ or $\mathbf{R}$ or $\mathbf{K}$ respectively. For a group of diffeomorphisms of a Hilbert manifold over $\mathbf{R}$ or over a non-Archimedean Banach manifold we have an analogous continuity of ρ_μ for a subgroup G'' of the entire group G (see [20, 22, 26, 28]).

By M_0 we denote a subspace in M , satisfying (a). Henceforth, we imply that R' contains all closed subsets from S belonging to $\mathbf{R}$, where G and $\mathbf{R}$ are complete.

For $\mu : Bf(S) \rightarrow G$ by $L(S, \mu, G)$ we denote the completion of a space of continuous $f : S \rightarrow G$ such that $\|f\|_d := \sup_{h \in C_b(S, G)} d(\int_S f(x)h(x) \mu(dx), 0) < \infty$ for each $d \in P''$, where $C_b(S, G)$ is a space of continuous

bounded functions $h : S \rightarrow G$. We suppose that for each sequence $(f_n : n) \subset L(S, \mu, G)$ for which $g \in L(S, \mu, G)$ exists with $d(f_n(x), 0) \leq d(g(x), 0)$ for every $d \in P''$, x and n , that f_n converges uniformly on each compact subset $V \subset S$ with $|\mu|(V) > 0$. In the cases $G = \mathbf{C}$ it coincides with $L^1(S, \mu, \mathbf{C})$ correspondingly, hence this supposition is the Lebesgue theorem. By $Y(v)$ we denote $L^1(\mathbf{K}, v, \mathbf{C})$.

Now we may define topologies and uniformities with the help of corresponding bases (see below) on $\mathbf{L} \subset G^{\mathbf{R}} \times R_o^{S' \times S} =: Y$ (or $P \subset G^{\mathbf{R}} \times R_o^{S' \times S} \times G^{S' \times K \times \mathbf{R}} =: Y$), $R_o \subset R \setminus \{0\}$. There are the natural projections $\pi : \mathbf{L} \rightarrow M_o$, $\pi(\mu, \rho_\mu(*, *) (\vee, \eta_\mu)) = \mu$, $\xi : \mathbf{L} \rightarrow R^{S' \times S}$, $\xi(\mu, \rho_\mu, (\vee \eta_\mu)) = \rho_\mu$, $\zeta : P \rightarrow G^{S' \times K \times \mathbf{R}}$, $\zeta(\mu, \rho_\mu, \eta_\mu) = \eta_\mu$. Let $\mathbf{H}$ be a filter on $\mathbf{L}$ or P , $U = U' \times U''$ or $U = U' \times U'' \times U'''$, U' and U'' be elements of uniformities on G , R and $Y(v)$ correspondingly, $\tau' \ni W \ni e$, $\tau \ni V \supset V_{U'} \ni e$, $V_{U'}$ is compact. By $[\mu]$ we denote (μ, ρ_μ) for $\mathbf{L}$ or $(\mu, \rho_\mu, \eta_\mu)$ for P , $\Omega := \mathbf{L} \vee P$, $[\mu](A, W, V) := [\mu^g(A), \rho_\mu(g, x), \vee \eta_{\mu^g}(t, U_*, A) \mid g \in W, x \in V, \vee t \in K]$. We consider $\mathbf{A} \subset \mathbf{R}$, then

$$(1) \mathbf{W}(\mathbf{A}, W, V_{U'}; U) := \{([\mu], [\nu]) \in \Omega^2 \mid ([\mu], [\nu])(A, W, V_{U'}) \subset U\};$$

$$(2) \mathbf{W}(\mathbf{S}; U) := \{([\mu], [\nu]) \in \Omega^2 \mid \{(B, g, x) : ([\mu], [\nu])(B, g, x) \in U\} \in \mathbf{S}\},$$

where $\mathbf{S}$ is a filter on $\mathbf{R} \times S' \times S^c$, S^c is a family of compact subsets $V' \ni e$.

$$(3) \mathbf{W}(\mathbf{F}, W, V; U) := \{([\mu], [\nu]) \in \Omega^2 \mid \{B : ([\mu], [\nu])(B, g, x) \in U, g \in W, x \in V\} \in \mathbf{F}\},$$

where $\mathbf{F}$ is a filter on $\mathbf{R}$ (compare with § 2.1 and 4.1[7]);

$$(4) \mathbf{W}(\mathbf{A}, \mathbf{G}; U) := \{([\mu], [\nu]) \in \Omega^2 \mid \{(g, x) : ([\mu], [\nu])(B, g, x) \in U, B \in \mathbf{A}\} \in \mathbf{G}\},$$

where $\mathbf{G}$ is a filter on $S' \times S^c$; suppose $\mathbf{U} \subset \mathbf{R} \times \tau'_e \times S^c$, Φ is a family of filters on $\mathbf{R} \times S' \times S^c$ or $\mathbf{R} \times S' \times S^c \times Y(v)$ (generated by products of filters $\Phi_{\mathbf{R}} \times \Phi_{S'} \times \Phi_{S^c}$ on the corresponding spaces), U' be a uniformity on (G, R) or $(G, R, Y(v))$, $\mathbf{F} \subset Y$. A family of finite intersections of sets $\mathbf{W}(A, U) \cap (\mathbf{F} \times \mathbf{F})$ (see (1)), where $(A, U) \in \mathbf{U} \times U'$ (or $\mathbf{W}(F, U) \cap (\mathbf{F} \times \mathbf{F})$ (see (2)), where $(F, U) \in (\Phi \times U')$ generate by the definition a base of uniformity of $\mathbf{U}$ -convergence (Φ -convergence respectively) on $\mathbf{F}$ and generate the corresponding topologies. For these uniformities are used notations

(i) $\mathbf{F}_U$ and $\mathbf{F}_\Phi$; $\mathbf{F}_{\mathbf{R} \times W \times V}$ is for $\mathbf{F}$ with the uniformity of uniform convergence

on $R \times W \times V$, where $W \in \tau'_e$, $V \in S^c$, analogously for the entire space Y ;

(ii) F_A denotes the uniformity (or topology) of pointwise convergence for

$A \subset R \times \tau'_e \times S^c =: Z$, for $A = Z$ we omit the index (see formula (1)). Henceforward, we use H' instead of H in 4.1.24[7], that is, $H'(A, \tilde{R})$ -filter on R generated by the base $[(L \in R : L \subset A \setminus K') : K' \in \tilde{R}, K' \subset A]$, where $\tilde{R} \subset R$ and $\tilde{R}$ is closed relative to the finite unions.

For example, let S be a locally $\mathbf{K}$ -convex space, S' be a dense subspace, G be a locally $\mathbf{L}$ -convex space, where $\mathbf{K}, \mathbf{L}$ are fields, $R = B(G)$ be a space of bounded linear operators on G , $R_o = GL(G)$ be a multiplicative group of invertible linear operators. Then others possibilities are: $S = X$ be a Banach space over $\mathbf{K}$, $S' = J_\mu$, $S'' = J_\mu^b$ as above; $S = G(t)$, $S' \supset S''$ are dense subgroups, $G = R$ be the field $\mathbf{R}$, M be an analytic Banach manifold over $\mathbf{K} \supset \mathbf{Q}_p$ (see [20]). The rest of the necessary standard definitions are recalled further when they are used.

5.2. Lemma. *Let R be a quasi- δ -ring with the weakest uniformity in which each $\mu \in M$ is uniformly continuous and $\Phi \subset \hat{\Phi}_C(R, S' \times S^c)$. Then $L(R, G, R, R')_\Phi$ (or $P(R, G, R, U_*)_\Phi$) is a topological space on which R_o acts continuously from the right.*

Proof. We recall that $\hat{\Phi} := \hat{\Phi}(X)$ denotes a family of filters F on X such that for each mapping $f : \Phi \rightarrow B(X)$ with $f(\Sigma) \subset \Sigma$ for each $\Sigma \in \Phi$ there exists a finite subset $\Psi \subset \Phi$ such that $\bigcup_{\Sigma \in \Psi} f(\Sigma) \in F$, where $B(X)$ denotes a family of all subsets in X , $\Phi_C(X) =: \Phi_C$ be a family of Cauchy filters on X . In view of proposition 4.2.2[7] the space of measures $M(R, G; R')_\Phi$ is a topological left- R -module. On the other hand, $\rho_{\lambda\mu}(a, x) = \lambda\rho_\mu(a, x)\lambda^{-1}$ for $\lambda \in R_o$, from the continuity of the inversion $\lambda \mapsto \lambda^{-1}$ in R_o and the multiplication in R for each entourage of the diagonal U'' in R there exists an entourage of a diagonal U''_1 in R and an open neighbourhood $U_4 \ni \lambda$ in R_o such that $\beta U''_1 \beta^{-1} \subset U''$ for each $\beta \in U_4$. Choosing $\Psi \subset \Phi$ for a given F we find $U''_1(\Sigma)$ and $U_4(\Sigma)$ for each $\Sigma \in \Psi$, then $\bigcap_{\Sigma \in \Psi} U''_1(\Sigma) =: \tilde{U}''$ and $\tilde{U}_4 = \bigcap_{\Sigma \in \Psi} U_4(\Sigma)$ are entourages of the diagonal and generate a neighbourhood of λ . This shows the continuity by λ for L and P , since for P is satisfied: $PD(b, \lambda f(x)) = \lambda PD(b, f(x))$ due to definition 4.1 for pseudo-differentiable f .

5.3. Proposition. (1). *Let T be a $\hat{\Phi}_4$ -filter on $M_o(R, G; R')$, $\{A_n\}$ be a disjoint $\Theta(R)$ -sequence, Σ be the elementary filter on R generated by*

$\{A_n : n \in \mathbf{N}\}$ and $\phi : \mathbf{M}_0 \times \mathbf{R} \rightarrow G$ with $\phi(\mu, A) = \mu(A)$. Then $\phi(\mathbf{T} \times \Sigma)$ converges to 0.

(2). Moreover, let $\mathbf{U}$ be a base of neighbourhoods of $e \in S'$, $\phi : \mathbf{L} \rightarrow G \times R$, $\phi(\mu, A, g) := (\mu^g(A), \rho_\mu(g, x))$, where $x \in A$. Then $(0, 1) \in \lim \phi(\mathbf{T} \times \Sigma \times \mathbf{U})$.

(3). If T is a Φ_4 -filter on $P(\mathbf{R}, G, R, U_*; R')$, $\psi(\mu, B, g, t, U_*) = [\mu(B); \rho_\mu(g, x); \eta_{\mu^g}(t, U_*, B)]$, then $(0, 1, 0) \in \lim \psi(\mathbf{T} \times \Sigma \times \mathbf{U})$ for each given $U_* \in S''$, where Σ and $\mathbf{U}$ as in (1,2).

Proof. We recall that for a set X , Θ -net in X is a net (I, f) in X such that for each increasing sequence $(i_n : n \in \mathbf{N}) \subset I$ is satisfied $(f(i_n) : n) \in \Theta$. A sequence $(x_n : n \in \mathbf{N}) \subset X$ is called a Θ -sequence, if $f : \mathbf{N} \rightarrow X$, $f(n) = x_n$, $(\mathbf{N}, f)$ is a Θ -net. Then $\Phi(\Theta, X) =: \Phi(\Theta)$ denotes a family of filters on X of the form $f(F)$, where (I, f) is a Θ -net in X , F is a filter of sections $[(\lambda \in I : \lambda \geq i) : i \in I]$ for a directed set I , $\hat{\Phi}(\Theta) := \Phi(\Theta)$. For a topological space X , $\Theta_j(X)$ denotes a family of sequences having converging subsequences for $j = 1$ or having limit points whilst $j = 2$. For a uniform space X , $\Theta_3(X)$ denotes a family of sequences having Cauchy subsequences $\Theta_4 := \Theta_2 \cup \Theta_3$, $\Phi_j(X) := \Phi(\Theta_j(X))$.

Then (1) follows from 4.2.6[7]. (2) From the restrictions (a,b) in 5.1 it follows that $\mu^g(A) = \int_A \rho_\mu(g, x) \mu(dx)$. If in addition $\lim \mu(F(A; R')) = \mu(A)$, then $\lim \mu(F(X \setminus A, R')) = \mu(X \setminus A)$, since by the definition 4.1.24[7] for each $\tau_G \ni D \ni 0$ for $A \in \mathbf{R}$ there exists $\tau_S \ni V \supset A$ with $\mu(V) - \mu(A) \in D$, where $F(A, R')$ is a filter generated by A and R' . Due to the Radonian property of μ and μ^g we get $\lim_{\mathbf{T} \times \Sigma} \mu^g(A) = 0$.

(3). Additionally to (1,2) it remains to verify that η_μ converges. In view of Theorems 4.4 and 4.7 (or §5.1) a pseudo-differential of order $b : \tilde{D}_{U_*}^b \mu = \nu$ is a measure for pseudo-differentiable μ . Due to (1) and the $\hat{\Phi}_4$ -condition we get $(0, 1, 0) \in \lim \psi(\mathbf{T} \times \Sigma \times \mathbf{U})$ for a given U_* , since in §§4.1 and 5.1 the integral is by $t \in \mathbf{K}$ for the Banach space X over $\mathbf{K}$ and by $B(\mathbf{K}, 0, 1)$ correspondingly for S that is not the Banach space.

5.4. Proposition. Let $\mathbf{H}$ be a $\hat{\Phi}_4$ -filter on $\mathbf{L}$ (or P) with the topology $\mathbf{F}$ (see §5.1.(ii)), $A \in \mathbf{R}$, $\tau_G \ni U \ni 0$, $\mathbf{H}'(A, R') \in \Psi_f(\mathbf{R})$. Then there are $L \in \mathbf{H}$, $\tilde{K} \in R'$ and an element of the uniformity $\mathbf{U}$ for $\mathbf{L}_{R'}$ or $P_{R'}$ such that $\tilde{\mathbf{K}} \subset A$, $L = [(\mu, \rho_\mu(g, x)) : M := \pi_{\mathbf{M}_0}(L) \ni \mu, \pi_{\tau'_e}(L) =: W \ni g$ (or $(\mu, \rho_\mu, \eta_\mu(*, *, U_*))$ and additionally $\tilde{D}_{U_*}^b \mu = PD(b, \eta_\mu)]$, $e \in W \in \tau'$, $\mu^g(B) - \nu^h(C) \in U$ (or in addition $(\tilde{D}_{U_*}^b \mu^g(B)) - (\tilde{D}_{U_*}^b \nu^h(C)) \in U$) for

$\tilde{K} \subset B \subset A$, $\tilde{K} \subset C \subset A$ for each $([\mu], [\nu]) \in \bar{\bar{L}}^2 \cap \mathcal{U}$, where $\bar{\bar{L}} := cl(L, \mathbf{L}_{R'})$ (or $cl(L, P_{R'})$), $\pi_{\mathbf{M}_0}$ is a projector from L into $\mathbf{M}_0$.

Proof. We recall that $\Psi(\mathbf{R}) := \hat{\Phi}(\Theta(\mathbf{R}))$, where $\Theta(\mathbf{R})$ is a family of sequences $(A_n : n \in \mathbf{N}) \subset \mathbf{R}$ for which there exists $\Omega \in \Sigma(\mathbf{R})$ with $(n \in \mathbf{N} : A_n \in \Omega)$ is infinite, $\Sigma(\mathbf{R}) := [(\bigcup_{n \in J} A_n : J \in \mathbf{B}(\mathbf{N})) : (A_n : n \in \mathbf{N}) \in \Gamma(\mathbf{R})]$, $\Gamma(\mathbf{R})$ is a family of disjoint sequences $(A_n : n \in \mathbf{N}) \subset \mathbf{R}$ for which $[\bigcup_{n \in J} A_n : J \subset \mathbf{N}] \subset \mathbf{R}$. A ring of sets Z is called a quasi- δ -ring, if each disjoint sequence $(A_n : n \in \mathbf{N})$ from $\mathbf{R}$ the union of which $\bigcup_n A_n = A$ is contained in a set $B \in \mathbf{R}$, $A \subset B$, has a subsequence $(A_{n_j} : j \in \mathbf{N}) \in \Gamma(\mathbf{R})$.

From Proposition 4.2.7[7] and the Radonian property of measures we have $\mu(B) - \nu(C) \in U'$ for each $(\mu, \nu) \in (\mathbf{M} \times \mathbf{M}) \cap \pi_{\mathbf{M}_0}(\mathcal{U})$. Then for each element D' of the uniformity on R there are $d \in P''$, $c > 0$ and L such that $\rho_\mu(g, x) - \rho_\nu(h, x') \in D'$ for $g, g' \in W$, $\mu, \nu \in M$, $d(x, x') < c$, since $H \in \hat{\Phi}_4$, where $\bar{\bar{M}} := cl(\mathbf{M}, E(\mathbf{R}, G; R')_{R'})$ is the closure of $\mathbf{M} \subset E(\mathbf{R}, G; R')$ in $E(\mathbf{R}, G; R')_{R'}$, $E(\mathbf{R}, G; R') := [\mu \in E(\mathbf{R}, G) : \mu \text{ is } R'\text{-regular}]$, that is, $\mu(F(A, R'))$ converges to $\mu(A)$ for each $A \in R$ (simply regular, if R' consists of closed subsets and conditions in definition 11.34[15] are satisfied). Then $F(A, R')$ is a filter on R generated by a base $[(B \in \mathbf{R} : \tilde{K} \subset B \subset A) : \tilde{K} \subset A, \tilde{K} \in R']$, $E(\mathbf{R}, G)$ is a set of exhaustive additive maps μ , that is, $\mu(A_n)$ converges to 0 for each $(A_n : n \in \mathbf{N}) \in \Gamma(\mathbf{R})$.

Then $\mu^g(B) - \mu(B) \in U'$, $\nu^h(C) - \nu(C) \in U'$ and for $3U' \subset U$ we get 5.4 for $\mathbf{L}$. From Theorems 4.4 and 4.7 (or §5.1), the Egorov conditions and the Lebesgue theorem we get 5.4 for P , since μ are probability measures and $\mathbf{L}_{R'}$ (or $P_{R'}$) correspond to uniformity from §5.1.(ii) with $A = R' \times \tau'_e \times S^c$. Indeed, $\mu^g(A) - \nu^h(A) = (\mu^g(A) - \mu^g(V_{U'})) + (\mu^g(V_{U'}) - \nu^h(V_{U'})) + (\nu^h(V_{U'}) - \nu^h(A))$, $\mu^g(A) = \int_A \rho_\mu(g, x) \mu(dx)$ for each $A \in Bf(S)$, for each $\tau_G \ni U' \ni 0$ there exists a compact subset $V'_U \subset A$ with $\mu^g(B) \in U'$ for each $B \in Bf(A \setminus V_{U'})$ and the same for ν^h (due to the Condition in 5.1 that R' contains $Bf(S)$). From the separability of S , S' and the equality of their topological weights to $\aleph_0$, restrictions 5.1(a,b) it follows that there exists a sequence of partitions $Z_n = [(x_m, A_m) : m, x_m \in A_m]$ for each $A \in Bf(S)$, $A_i \cap A_j = \emptyset$ for each $i \neq j$, $\bigcup_m A_m = A$, $A_m \in Bf(S)$, such that $\lim_{n \rightarrow \infty} (\mu^g(A) - \sum_j \rho_\mu(g, x_j) \mu(A_j)) = 0$ and the same for ν , moreover, for $V_{U'}$ each Z_n may be chosen finite. Then there exists $W \in \tau'_e$ with $W \times (S \setminus V^2) \subset (S \setminus V)$, $\tau_e \ni V \subset V^2$, $\nu^g(B)$ and $\mu^g(B) \in U'$ for each $B \in Bf(S \setminus V^2)$ (for $G = \mathbf{R}$) and $g \in W$ (see 5.1.(a)). Then from $A = [A \cap (S \setminus V^2)] \cup [A \cap V^2]$ and the existence of

compact $V'_{U'} \subset V$ with $\mu(E) \in U'$ for each $E \in Bf(V \setminus V'_{U'})$ and the same for ν , moreover, $(V'_{U'})^2$ is also compact, it follows that $\mu^g(B) - \nu^h(C) \in U$ for $9U' \subset U$, since $R' \supset Bf(S)$, where W satisfies the following condition $\mu^g(V'_{U'}) - \nu^h(V'_{U'}) \in U'$ for $V'_{U'} \subset V^2$ due to 5.1.(b), $\mu(B) - \nu(C) \in U'$, $WV'_{U'} \subset (V'_{U'})^2$ due to precompactness of W in S . Since pseudo-differentiable measures are also quasi-invariant, hence for them Proposition 5.4 is true.

Now let $[\mu] \in \lim H$, $A \in Bf(S)$, then $\eta_\mu \in \lim \zeta(H)$ in $Y(v)$ and there exists a sequence η_{μ_n} such that $\int_{\mathbf{K}} \eta_{\mu_n}(\lambda, U_*, A) v(d\lambda) = \tilde{D}_{U_*}^b \mu_n(A)$ due to §§4.4, 4.7 or 5.1 and $\lim_{n \rightarrow \infty} \tilde{D}_{U_*}^b \mu_n(A) = \int_{\mathbf{K}} \eta_\mu(\lambda, U_*, A) v(d\lambda) =: \kappa(A)$ due to the Lebesgue theorem. From $\eta_\mu(\lambda, U_*, A \cup B) = \eta(\lambda, U_*, A) + \eta(\lambda, U_*, B)$ for $A \cap B = \emptyset$, $B \in Bf(S)$ and the Nikodym theorem [7] it follows that $\nu(A)$ is the measure on $Bf(S)$, moreover, $\kappa(A) = \tilde{D}_{U_*}^b \mu(A)$. Since $\mu^g(A) = \int_A \rho_\mu(g, x) \mu(dx)$ for $A \in Bf(S)$ for $g \in S'$, then

$$\begin{aligned} \eta_{\mu^g}(\lambda, U_*, A) &= j(\lambda) g(\lambda, 0, b) [\mu^g(A) - \mu^g(U_\lambda^{-1} A)] \\ &= j(\lambda) g(\lambda, 0, b) \int_A \rho_\mu(g, x) [\mu(dx) - \mu^{U_\lambda}(dx)] \end{aligned}$$

and in view of the Fubini theorem there exists

$$\tilde{D}_{U_*}^b \mu^g(A) = \int_A \left[\int_K \rho_\mu(g, x) j(\lambda) g(\lambda, 0, b) [\mu(dx) - \mu^{U_\lambda}(dx)] v(d\lambda) \right],$$

where $j(t) = 1$ for $S = X$ and $j(t)$ is the characteristic function of $B(\mathbf{K}, 0, 1)$ for S that is not the Banach space X . Then μ -a.e. $\tilde{D}_{U_*}^b \mu^g(dx) / \tilde{D}_{U_*}^b \mu(dx)$ coincides with $\rho_\mu(g, x)$ due to 5.1.(a,b), hence, $(\tilde{D}_{U_*}^b \mu^g, \rho_{\mu^g})$ generate the Φ_4 -filter in L arising from the $\hat{\Phi}_4$ -filter in P . Then we estimate $\tilde{D}_{U_*}^b (\mu^g - \nu^h)(A)$ as above $\mu^g(A) - \nu^h(A)$. Therefore, we find for the Φ_4 -filter corresponding L , since there exists $\delta > 0$ such that $U_\lambda \in W$ for each $|\lambda| < \delta$. For Φ_4 -filter we use the corresponding finite intersections $W_1 \cap \dots \cap W_n = W$, where W_j correspond to the Φ_4 -filters H_j .

5.5. Corollary. *If $\{H'(A, R') : A \in R\} \subset \Psi_f(R)$, T is a $\hat{\Phi}_4$ -filter in L , U is an element of uniformity in L (or P), then there are $L \in T$ and an element V of the uniformity in $L_{R'}$ (or $P_{R'}$) such that $\bar{L}^2 \cap V \subset \bar{L}^2 \cap U$.*

Proof. In view of Corollary 4.2.8[7] and Theorem 5.4 for $U_M = \pi_M \times \pi_M U$ there exists an entourage of the diagonal, V_M in $E(R, G; R')_{R'}$, such that $\bar{M}^2 \cap V_M \subset \bar{M}^2 \cap U_M$, where $\pi_M : L \rightarrow M$ (or $P \rightarrow M$) is a projector, $M(R, G)$ is a set of measures on R with values in G , $M(R, G; R') := [\mu \in M(R, G) :$

μ is R' -regular]. Since $M(R, G; R')_{R'} \subset E(R, G; R')_{R'}$, then there exists an entourage of the diagonal, V in $L_{R'}$ (or $P_{R'}$), such that $\pi_M \times \pi_M V \subset V_M \cap \bar{M}^2$.

5.6. Lemma. *Let R be a quasi- σ -ring directed by the inclusion, U be an upper directed subset in R , H be a $\hat{\Phi}_4$ -filter in L (or P), $0 \in U \ni \tau_G$. Then there are $M \in H$, $A \in U$, an element of the uniformity U' in $L_{R'}$ (or $P_{R'}$) such that $\mu^g(B) - \nu^h(C) \in U$ (or in addition $\tilde{D}_{U*}^b \mu^g(B) - \tilde{D}_{U*}^b \nu^h(C) \in U$) for each $([\mu], [\nu]) \in \bar{L}^2 \cap U'$, $g, h \in W$, W is defined by a projection of U' onto S' , $B, C \in U$ with $A \leq B$, $A \leq C$.*

Proof. For each C and $E \in U$ by the definition there are $F \in U$ with $C \leq F$ and $E \leq F$. There are open subsets P and V in S and $W \in \tau'_e$ with $WP \subset P^2 \subset V$ such that $(\mu - \nu)(B) \in U$ for each $B \subset S \setminus P$ and $(\mu, \nu) \in \bar{M}^2 \cap \pi_{M_0}(U')$. Indeed, $\pi_{M_0}H$ is a base of a filter T in M_0 . Therefore, $g(B \cap S \setminus P^2) \subset S \setminus P$ for $g \in W$, consequently, $[\mu^g - \nu^g](B \cap S \setminus P^2) \in U$. In view of §5.1.(a,b) for D, V there are $W, c > 0, d \in P''$ such that $\rho_\mu(g, x) - \rho_\nu(h, x') \in D$ for $g, h \in W$ and $d(x, x') < c, x, x' \in V_c$, consequently, $\mu^g(B) - \nu^h(C) = [\mu^g(B \cap P^2) - \nu^h(C \cap P^2)] + [\mu^g(B \setminus P^2) - \nu^h(C \setminus P^2)] \in 3(DU + U)$. Modifying the proof of §4.2.9 [7] we get the statement of this lemma for L . For P an estimate of $\tilde{D}_{U*}^b \mu^g(B) - \tilde{D}_{U*}^b \nu^h(C)$ may be done analogously to the proof of Proposition 5.4.

5.7. Theorem. *Let H be a $\hat{\Phi}_4$ -filter in L (or P), $\{A_n : n \in \mathbb{N}\} \in \Gamma(R)$, $\tau_G \ni U \ni 0$. Then there are $L \in H$, $M \in B(\mathbb{N})$ and an element U of the uniformity in L such that $\mu^g(\cup(A_n : n \in M')) - \nu^h(\cup(A_n : n \in M'')) \in U$ (or in addition $(\tilde{D}_{U*}^b \mu^g(\cup_{n \in M'} A_n) - \tilde{D}_{U*}^b \nu^h(\cup_{n \in M''} A_n)) \in U$ for each $([\mu], [\nu]) \in \bar{L}^2 \cap U$ and $M', M'' \in B(\mathbb{N})$ with $M \cap M' = M \cap M''$. If $\{H'(A, R')\} \subset \Psi_f(R)$, then U may be chosen as an element of the uniformity in $L_{R'}$ (or $P_{R'}$), where $\bar{L} := cl(L, L)$ (or $\bar{L} := cl(L, P)$).*

Proof. We recall that $B_f(\mathbb{N})$ denotes the family of finite subsets in $\mathbb{N}$, $\Psi_f(R) := \hat{\Phi}(\Theta_f(R))$, $\Theta_f(R)$ is a family of sequences $(A_n : n \in \mathbb{N}) \subset R$ for which there exists $\Omega \in \Sigma_f(R)$ with $card[n \in \mathbb{N} : A_n \in \Omega] = \aleph_0$, $\Sigma_f(R) := [(\cup_{n \in M} A_n : M \in B_f(\mathbb{N})) : (A_n : n \in \mathbb{N}) \in \Gamma(R)]$. Let $\phi(M) := \cup\{A_n : n \in M\}$, $M \subset B_f(\mathbb{N})$, Σ be a filter of neighbourhoods of $\emptyset$ in $B_f(\mathbb{N})$, then $\phi(\Sigma) \in \Psi_f(R)$. In view of §4.1.14 [7] we have $\mu(\phi(\Sigma)) \rightarrow 0$ for each $\mu \in M(R, G, R')$. We take a symmetric neighbourhood $V' \ni 0$ in G with $3V' \subset U$. Then there are $L \in H$ and $M \in B_f(\mathbb{N})$ such that $\mu^g(\phi(M')) - \nu^h(\phi(M'')) \in V'$ (or in addition $\tilde{D}_{U*}^b \mu^g(\phi(M')) - \tilde{D}_{U*}^b \nu^h(\phi(M'')) \in V'$) for each $([\mu], [\nu]) \in \bar{L}^2 \cap U$ and $M', M'' \in B(\mathbb{N} \setminus M)$ (see Lemma 5.6). We choose U such that $\mu^g(\phi(M_o)) -$

$\nu^h(\phi(M_o)) \in V'$ (or in addition $\tilde{D}_{U_*}^b \mu^g(\phi(M_o)) - \tilde{D}_{U_*}^b \nu^h(\phi(M_o)) \in V'$) for each $([\mu], [\nu]) \in \mathbf{U}$ and $M_o \subset M$, consequently, $\mu^g(\phi(M')) - \nu^h(\phi(M'')) \in 3V' \subset U$ (or additionally $\tilde{D}_{U_*}^b \mu^g(\phi(M')) - \tilde{D}_{U_*}^b \nu^h(\phi(M'')) \in 3V'$) for $M \cap M' = M \cap M''$, the last statement follows from Corollary 5.5.

5.8. Corollary. *Let $(A_n : n \in \mathbf{N}) \in \Gamma(\mathbf{R})$, $\mathbf{H}$ be a $\hat{\Phi}_4$ -filter in $\mathbf{L}$ (or P), $\tau_G \ni U \ni 0$, then there are $(L, M) \in \mathbf{H} \times \mathbf{B}_f(\mathbf{N})$ such that $\mu^g(\cup(A_n : n \in M')) \in U$ (or additionally $\tilde{D}_{U_*}^b \mu^g(\cup_{n \in M'} A_n) \in U$) for $[\mu] \in \bar{L}$ and $M' \in \mathbf{B}(\mathbf{N} \setminus M)$.*

Proof. Let us take a fixed ν and W from §5.7 with $\nu(\cup_{n \in M'} A_n) \in \tilde{U}$ (or additionally $\tilde{D}_{U_*}^b \nu(\cup_{n \in M'} A_n) \in \tilde{U}$) and $(\mu^g - \nu)(\cup_{n \in M'} A_n) \in \tilde{U}$, where $2\tilde{U} \subset U$, $0 \in \tilde{U} \in \tau_G$, $g \in W$.

5.9. Theorem. *Let $\mathbf{U} \subset \mathbf{R} \times \tau'_e \times S^c =: Z$ be such that $\text{id} : \mathbf{L}_{\mathbf{U}} \rightarrow \mathbf{L}$ (or $\text{id} : P_{\mathbf{U}} \rightarrow P$) be uniformly Φ_4 -continuous, $\{\mathbf{H}'(A, R') : A \in \mathbf{R}\} \subset \Psi_f(\mathbf{R})$. Then*

(a) $\mathbf{L}$ (or P) is Φ_4 -closed in $G^{\mathbf{R}} \times R_o^{S' \times S}$ (or $G^{\mathbf{R}} \times R_o^{S' \times S} \times G^{S' \times K \times \mathbf{R}}$), $\mathbf{M}_o$ is Φ_4 -closed in $G^{\mathbf{R}}$;

(b) if G and R are Φ_i -compact, then $\mathbf{L}_{\mathbf{U}}$ (or $P_{\mathbf{U}}$) is Φ_i -compact, where $i \in (1, 2, 3, 4)$;

(c) if $G \times R_o$ is sequentially complete, then $\mathbf{L}_{\mathbf{U}}$ (or $P_{\mathbf{U}}$) is also;

(d) if $(0, 0)$ is the G_δ -subset in $G \times R$, then $\mathbf{L}_{\mathbf{U}}$ (or $P_{\mathbf{U}}$) is Φ_2 -compact; if additionally $G \times R$ is sequentially complete, then $\mathbf{L}_{\mathbf{U}}$ (or $P_{\mathbf{U}}$) is Φ_4 -compact;

(e) $\mathbf{L}_{\mathbf{U}}$ (or $P_{\mathbf{U}}$) is Hausdorff.

Proof. We recall that a subset $A \subset E$ of a topological or of a uniform space E is called Φ_i -closed (or compact), if for each $\Phi_i(A)$ -filter F is satisfied $\lim F \subset A$ (or $\lim F \neq \emptyset$ respectively, that is, this definition of compactness differs from the usual). In view of Theorem 4.2.14[7] and Proposition 5.4 is accomplished 5.9(a).

From §§2.1.14, 1.8.11[7] and Theorem 5.7, Φ_i -compactness of $\mathbf{M}$ and $\mathbf{M}_o$, also from the completeness of R_o , $L^1(K, v, \mathbf{C})$ it follows (b).

Then (c) follows from (a) and §1.8.7[7];

(d) follows from (c) and §1.6.4[7], since $G \times R$ is Φ_2 -compact. From $\mathbf{M}(\mathbf{R}, G; R')_{\mathbf{U}}$ (see §4.2.14[7]), $R_o^{S' \times S}$ and $L^1(K, v, \mathbf{C})$ being Hausdorff it follows (d).

5.10. Theorem. *Let Φ be a set of $\hat{\Phi}_4$ -filters on $\mathbf{L}$ (or P), also let $\phi : \mathbf{L} \times \mathbf{R} \times S' \times S(\vee \times \mathbf{K}) \rightarrow G \times R_o(\vee \times Y(v))$ be such that $(\phi(\mu, \rho_\mu(*, *) (\vee \eta_\mu), A, g, x) := (\mu(A), \rho_\mu(g, x) (\vee \eta_{\mu^g}(t, U_*, A)))$. If $\{\mathbf{H}'(A, R') | A \in \mathbf{R}\} \subset \Psi_f(\mathbf{R})$, then a map-*

ping $\phi : \mathbf{R} \times S' \times S(\mathbf{V} \times \mathbf{K}) \rightarrow G_{\Phi'}^{\mathbf{M}_0(\mathbf{R}), G; R'} \times R_o^{S' \times S}(\mathbf{V} \times Y(v))$ gives R' -regular quasi-invariant measure (or in addition a pseudo-differentiable measure) $(A, g, x) \rightarrow \phi(*, A, g, x)$ (or $(A, g, x, t) \rightarrow \phi(*, A, g, x, t)$), satisfying Conditions 5.1.(a,b), where $\Phi' := \pi_{\mathbf{M}_0}(\Phi)$, in $R_o^{S' \times S}$ the topology corresponds to the topology in $\mathbf{L}$.

Proof follows from §§5.4 and 5.8.

5.11. Note. Let it be a sequence $\{[\mu_n] : n \in \mathbf{N}\}$ of quasi-invariant measures (or pseudo-differentiable measures) converging uniformly in uniformity 5.1.(ii) and fulfilling Conditions 5.1.(a,b) then in accordance with Corollary 5.8 $\{(\mu_n)^g : n \in \mathbf{N}\}$ (or also $[\tilde{D}_{U*}^b \mu_n^g : n]$) is uniformly σ -additive for each fixed $g \in S'$. Moreover, it is uniformly σ -additive by $g \in W$ for each given $B \in \mathbf{R}$ such that $gB \subset V$ for suitable open W in S' and V in S . For $\mathbf{L}$ this means that for each $0 \in D \in \tau_R$ and $e \in U \in \tau_G$ there are $W \ni e$, $d \in P''$, $c > 0$, n and $V \ni e$, a compact subset V_U , $e \in V_U \subset V$ with $\mu_m(C) \in U$ (or $\tilde{D}_{U*}^b \mu_m(C) \in U$ in addition for P) for $C \in \mathbf{R}$ and $C \subset S \setminus V_U$, with $\rho_{\mu_m}(g, x) - \rho_{\mu_j}(h, x') \in D$ (or in addition $\tilde{D}_{U*}^b (\mu_m^g - \mu_j^h)(A) \in U$ for $A \in Bf(V_U)$ for P) whilst $g, h \in W$ for each $x, x' \in V_U$ with $d(x, x') < c$ and $m, j > n$. In view of Theorem 5.9 there exists $\lim_{n \rightarrow \infty} (\mu_n, \rho_{\mu_n}(g, x)) = (y, d(y; g, x)) \in \mathbf{L}$, that is, a quasi-invariant measure (or pseudo-differentiable measure $\lim_n [\mu]_n = [\mu] \in P$). Therefore, they are analogs of the Radon theorem for quasi-invariant measure and pseudo-differentiable measures.

6 Appendix.

Suppose $X = c_0(\omega_0, \mathbf{K})$ is a Banach space over a locally compact non-Archimedean infinite field $\mathbf{K}$ with a non-trivial valuation and I is a unit operator on X . If A is an operator on X , then in some basis of X we have an infinite matrix $(A_{i,j})_{i,j \in \mathbf{N}}$, so we can consider its transposed matrix A^t . If in some basis the following equality is satisfied $A^t = A$, then A is called symmetric.

A.1. Lemma. *Let $A : X \rightarrow X$ be a linear invertible operator with a compact operator $(A - I)$. Then there exist an orthonormal basis $(e_j : j \in \mathbf{N})$ in X , invertible linear operators $C, E, D : X \rightarrow X$ with compact operators $(C - I)$, $(E - I)$, $(D - I)$ such that $A = SCDE$, D is diagonal, C is lower triangular and E is upper triangular, S is an operator transposing a finite number of vectors from an orthonormal basis in X . Moreover, there exists*

$n \in \mathbf{N}$ and invertible linear operators $A', A'' : X \rightarrow X$ with compact operators $(A' - I)$, $(A'' - I)$ and $(A'_{i,j} - \delta_{i,j} = 0)$ for i or $j > n$, A'' is an isometry and there exist their determinants $\det(A')\det(A'') = \det(A)$, $|\det(A'')|_{\mathbf{K}} = 1$, $\det(D) = \det(A)$. If in addition A is symmetric, then $C^t = E$ and $S = I$.

Proof. In view of Lemma 2.2[36] for each $c > 0$ there exists the following decomposition $X = Y \oplus Z$ into $\mathbf{K}$ -linear spaces such that $\|(A - I)|_Z\| < c$, where $\dim_{\mathbf{K}} Y = m < \aleph_0$. In the orthonormal basis $(e_j : j)$ for which $\text{sp}_{\mathbf{K}}(e_1, \dots, e_m) = Y$ for $c \leq 1/p$ we get $A = A'A''$ with $(A - I)|_Z = 0$, $|A''_{i,j} - \delta_{i,j}| \leq c$ for each i, j such that $(A'_{i,j} - \delta_{i,j}) = 0$ for i or $j > n$, where $n \geq m$ is chosen such that $|A_{i,j} - \delta_{i,j}| \leq c^2$ for $i > n$ and $j = 1, \dots, m$, $A_{i,j} := e_i^*(Ae_j)$, e_i^* are vectors e_i considered as linear continuous functionals $e_i^* \in X^*$. Indeed, $(A_{i,j} : i \in \mathbf{N}) = Ae_j \in X$ and $\lim_{i \rightarrow \infty} A_{i,j} = 0$ for each j . From the form of A'' it follows that $\|A''e_j - e_j\| \leq 1/p$ for each j , consequently, $\|A''x\| = \|x\|$ for each $x \in X$. Since $A'' = (A')^{-1}A$, $(A - I)$ and $(A' - I)$ being compact, hence $(A'' - I)$ is compact together with $(A^{-1} - I)$, $((A')^{-1} - I)$ and $((A'')^{-1} - I)$. Moreover, there exists $\lim_{k \rightarrow \infty} \det(A)_k = \det(A) = \lim_k \det((A')_k(A'')_k) = \lim_k \det(A')_k \det(A'')_k = \det(A')\det(A'')$, where $(A)_k := (A_{i,j} : i, j \leq k)$. This follows from the decompositions $X = Y_k \oplus Z_k$ for $c = c(k) \rightarrow 0$ whilst $k \rightarrow \infty$. This means that for each $c(k) = p^{-k}$ there exists $n(k)$ such that $|A_{i,j} - \delta_{i,j}| < c(k)$, $|A'_{i,j} - \delta_{i,j}| < c(k)$ and $|A''_{i,j} - \delta_{i,j}| < c(k)$ for each i or $j > n(k)$, consequently, $|A \begin{pmatrix} 1 \dots n(k) \\ 1 \dots n(k) \end{pmatrix}_{i_1 \dots i_q} - A \begin{pmatrix} 1 \dots n(k) \\ 1 \dots n(k) \end{pmatrix}_{\delta_{i_1, j_1} \dots \delta_{i_q, j_q}}| < c(k)$, where $A \begin{pmatrix} i_1 \dots i_r \\ j_1 \dots j_r \end{pmatrix}$ is a minor corresponding to rows $i_1, \dots, i_r$ and columns $j_1, \dots, j_r$ for $r, q \in \mathbf{N}$. From the ultrametric inequality it follows that $|\det(A'') - 1| \leq 1/p$, hence $|\det(A'')|_{\mathbf{K}} = 1$, $\det(A'')_k \neq 0$ for each k , $\det(A')_k = \det(A')_n$ for each $k \geq n$. Using the decomposition of $\det(A')_n$ by the last row (analogously by the column) we get $A'_{n,j} \neq 0$ and a minor $A' \begin{pmatrix} 1 \dots n-1 \\ 1 \dots j-1, j+1, n \end{pmatrix} \neq 0$. Permuting the columns j and n (or rows) we get as a result a matrix $(\bar{A})_n$ with $\bar{A} \begin{pmatrix} 1 \dots n-1 \\ 1 \dots n-1 \end{pmatrix} \neq 0$. Therefore, by the denumeration of the basic vectors we get $A' \begin{pmatrix} 1 \dots k \\ 1 \dots k \end{pmatrix} \neq 0$ for each $k = 1, \dots, n$, since $|\det(\bar{A})_n| = |\det(A')_n|$.

Therefore, there exists the orthonormal basis $(e_j : j)$ such that $A \begin{pmatrix} 1 \dots j \\ 1 \dots j \end{pmatrix} \neq 0$ for each j and $\lim_j A \begin{pmatrix} 1 \dots j \\ 1 \dots j \end{pmatrix} = \det(A) \neq 0$. Applying to $(A)_j$ the Gaussian decomposition and using compactness of $A - I$ due to formula (44) in §II.4[12], which is valid in the case of $\mathbf{K}$ also, we get $D = \text{diag}(D_j : j \in \mathbf{N})$, $D_j = A \begin{pmatrix} 1 \dots j \\ 1 \dots j \end{pmatrix} / A \begin{pmatrix} 1 \dots j-1 \\ 1 \dots j-1 \end{pmatrix}$; $C_{g,k} = A \begin{pmatrix} 1 \dots, k-1, g \\ 1 \dots, k-1, k \end{pmatrix} / A \begin{pmatrix} 1 \dots k \\ 1 \dots k \end{pmatrix}$; $E_{k,g} = A \begin{pmatrix} 1 \dots, k-1, k \\ 1 \dots, k-1, g \end{pmatrix} / A \begin{pmatrix} 1 \dots k \\ 1 \dots k \end{pmatrix}$

for $g = k + 1, k + 2, \dots, k \in \mathbf{N}$. Therefore, $(C - I)$, $(D - I)$, $(E - I)$ are the compact operators, $C_{i,j}$, D_j , $E_{i,j} \in \mathbf{K}$ for each i, j . Particularly, for $A^t = A$ (A^t denotes the transposed matrix for A) we get $E_{k,g} = C_{g,k}$.

A.2. Notes. 1. An isometry operator S does not influence on the results given above by the measures, since $A|Z$ and $S^{-1}A$ on X have decompositions of the form CDE , where $X \ominus Z$ is a finite-dimensional subspace of X over $\mathbf{K}$, hence $X \ominus Z$ is locally compact for the locally compact field $\mathbf{K}$. Therefore, on $X \ominus Z$ there exists the Haar measure with values in $\mathbf{R}$ or $\mathbf{Q}_s$ with $s \neq p$, $\mathbf{K} \supset \mathbf{Q}_p$.

A.2.2. For compact $A - I$ we can construct the following decomposition $A = BDB^tC$, where $BB^t = I$, $CC^t = I$, D is diagonal, B , D and E are operators on $c_0(\omega_0, \mathbf{C}_p)$ with $\lim_{i,j \rightarrow \infty} (B_{i,j} - \delta_{i,j}) = \lim_j (D_j - 1) = 0$, $\lim_{i,j \rightarrow \infty} (C_{i,j} - \delta_{i,j}) = 0$. But, in general, matrix elements of B, C, D are in $\mathbf{C}_p$ and may be in $\mathbf{C}_p \setminus \mathbf{K}$, since from the secular equation $\det(A - \lambda I)$ even for symmetric matrix A over $\mathbf{K}$ in general may appear $p^{1/n}$ for $n \in \mathbf{N}$, but $\mathbf{R}$ is not contained in $\mathbf{K}$ and $\mathbf{C}_p$ is not locally compact. This decomposition is not used in the present article for the construction of quasi-invariant measures on a Banach space X over $\mathbf{K}$, since on $\mathbf{C}_p$ there is not any non-trivial invariant measure (or even quasi-invariant relative to all shifts from $\mathbf{C}_p$) and may be $B_{i,j}$, D_j , $C_{i,j} \in \mathbf{C}_p \setminus \mathbf{K}$. Instead of it (and apart from [38]) we use the decomposition given in the Lemma A.1.

A.3. Remarks. Let $\mathbf{K}$ be a non-Archimedean infinite field with a non-trivial valuation. Let X and Y be normed spaces over $\mathbf{K}$, and $F : U \rightarrow Y$ be a function, where $U \subset X$ is an open subset. The function F is called differentiable, if for each $t \in \mathbf{K}$, $x \in U$ and $h \in X$ such that $x + th \in U$ there exists $DF(x, h) := \{dF(x + th)/dt \mid t = 0\} = \lim_{t \rightarrow 0, t \neq 0} \{F(x + th) - F(x)\}/t$ and $DF(x, h)$ linear by h , that is, $DF(x, h) =: F'(x)h$, where $F'(x)$ is a bounded linear operator (derivative). Let $\Phi^1 F(x; h; t) := \{F(x + th) - F(x)\}/t$ for each $t \neq 0$, $x \in U$, $x + th \in U$, $h \in X$. If this function $\Phi^1 F(x; h; t)$ has a continuous extension $\bar{\Phi}^1 F$ on $U \times V \times S$, where U and V are open neighbourhoods of x and 0 in X , $S = B(\mathbf{K}, 0, 1)$ and $\|\bar{\Phi}^1 F(x; h; t)\| := \sup\{\|\Phi^1 F(x; h; t)\|/\|h\| : x \in U, 0 \neq h \in V, t \in S\} < \infty$ and $\bar{\Phi}^1 F(x; h; 0) = F'(x)h$, then F is called continuously differentiable on U , the space of all such F is denoted by $C^1(U, Y)$, where $B(X, y, r) := \{z \in X : \|z - y\|_X \leq r\}$. By induction we define $\Phi^{n+1} F(x; h(1), \dots, h(n+1); t(1), \dots, t(n+1)) := \{\Phi^n F(x + t(n+1)h(n+1); h(1), \dots, h(n); t(1), \dots, t(n)) - \Phi^n F(x; h(1), \dots, h(n); t(1), \dots, t(n))\}/t(n+1)$ and the space $C^n(U, Y)$ [22, 24].

The family of all bijective surjective mappings of U onto itself of class C^n is called the diffeomorphism group and it is denoted by $\text{Diff}^n(U) := C^n(U, U) \cap \text{Hom}(U)$.

A.4. Theorem. *Let $\mathbf{K}$ be a spherically complete non-Archimedean infinite field with non-trivial valuation, $F \in C^n(U, Y)$ with $n \in \mathbf{N} := \{1, 2, \dots\}$, $F : X \rightarrow Y$, X and Y be Banach spaces, U be an open neighbourhood of $y \in X$, $F(X) = Y$ and $[F'(y)]^{-1}$ be continuous, where $n \in \mathbf{Z}$, $0 \leq n < p$ when $\text{char}(\mathbf{K}) = p > 0$, $n \geq 0$ when $\text{char}(\mathbf{K}) = 0$. Then there exists $r > 0$ and a locally inverse operator $G = F^{-1} : V := B(Y, F(y), \|F'(y)\|r) \rightarrow B(X, y, r)$, moreover, $G \in C^n(V, X)$ and $G'(F(y)) = [F'(y)]^{-1}$.*

Proof. Let us consider an open subset $U \subset X$, $[y, x] := [z : z = y + t(x - y), |t| \leq 1, t \in \mathbf{K}] \subset U$ and $g : Y \rightarrow \mathbf{K}$ be a continuous $\mathbf{K}$ -linear functional, $f(t) := g(F(y + t(x - y)))$. Such family of non-trivial g separating points of X exists due to the Hahn-Banach theorem [33, 34], since $\mathbf{K}$ is spherically complete. Then there exists $f'(t) = g(F'(y + t(x - y))(x - y))$, $f \in C^n(B(\mathbf{K}, 0, 1), \mathbf{K})$, consequently, for $F'(z) \neq 0$, $z \in [x, y]$ there exists $b > 0$ such that for $g(F'(z)) \neq 0$ we have $|g(F(x + t(x - y))) - g(F(x))| = |g(F'(z)(x - y))| \times |t|$ for each $|t| < b$. From $|g(u(z)) - g(u(x))| = |g(z) - g(x)|$ and the Hahn-Banach it follows, that u is the local isometry, where $u(z) := [F'(y)]^{-1}(F(z))$, $z, x \in B(X, y, r)$, r is chosen such that $\|(\bar{\Phi}^1 u(x, h, t)) - h\| < \|h\|/2$ for each $x \in B(X, y, r)$, $h \in B(X, 0, r)$, $t \in B(\mathbf{K}, 0, 1)$. Therefore, $F(B(X, y, r)) \subset B(Y, F(y), \|F'(y)\|r) =: S$. Applying to the function $H(z) := z - [F'(y)]^{-1}(F(z) - q)$ for $q \in S$ the fixed point theorem we get z' such that $F'(z') = q$, since $\|H(x) - H(z)\| < b \times \|x - z\|$. From the uniform continuity by (g, h) differentiability $f_{g,h}(t) := g(F(y + th))$ by t it follows continuous differentiability of G .

A.5. Theorem. *Let $f \in C^n(U, Y)$, where $n \in \mathbf{N}$, Y is a Banach space, $n \in \mathbf{Z}$, $0 \leq n < p$ when $p := \text{char}(\mathbf{K}) > 0$, $n \geq 0$ when $\text{char}(\mathbf{K}) = 0$. Then for each x and $y \in U$ is accomplished the following formula: $f(x) = f(y) + \sum_{j=1}^{n-1} f^{(j)}(y)(x - y)^j/j! + R_n(x, y)(x - y)^{n-1}$, where $f^{(j)}(y)(x - y)^j = (\bar{\Phi}^j f)(y; x - y, \dots, x - y; 0, \dots, 0) \times j!$, $f^{(j)}(y)h^j := f^{(j)}(y)(h, \dots, h)$, $f^{(j)}(y) : U \rightarrow L_j(X^{\otimes j}, Y)$, $R_n(x, y) : U^2 \rightarrow L_{n-1}(X^{\otimes(n-1)}, Y)$ with $R_n(x, y) = o(\|x - y\|)$, where $L_j(X^{\otimes j}, Y)$ is the Banach space of continuous polylinear operators from $X^{\otimes j}$ to Y , U is open in X .*

Proof. For $n = 1$ this formula follows from the definition of $\bar{\Phi}^1 f$. For $n = 2$ let us take $R_2(x, y) = \bar{\Phi}^2 f(x; y, y)$. Evidently, $C^{n-1}(U, Y) \supset C^n(U, Y)$.

Let the statement be true for $n - 1$, where $n \geq 3$. Then from $\bar{\Phi}^{n-1}f(y + t_n(x - y); x - y, \dots, x - y; t_1, \dots, t_{n-1}) = \bar{\Phi}^{n-1}f(y; x - y, \dots, x - y; t_1, \dots, t_{n-1}) + (\bar{\Phi}^1(\bar{\Phi}^{n-1}f(y; x - y, \dots, x - y; t_1, \dots, t_{n-1}))(y; x - y; t_n)) \times t_n$ and continuity of $\bar{\Phi}^n$ it follows that $R_n(x, y)(x - y)^{n-1} = \bar{\Phi}^n f(y; x - y, \dots, x - y; 0, \dots, 0, 1) - f^{(n)}(y)(x - y)^n/n! = \alpha(x - y)(x - y)^{n-1}$, where $\lim_{x \rightarrow y, x \neq y} \alpha(x - y)/\|x - y\| = 0$, that is, $\alpha(x - y) = o(\|x - y\|)$.

References

- [1] I.Ya. Aref'eva, B. Dragovich, I.V. Volovich. "On the p -adic summability of the anharmonic oscillator" // Phys. Lett., **B 200** (1988), 512-514.
- [2] A. H. Bikulov, I.V. Volovich. " p -Adic Brounian motion" // Izvest. Russ. Acad. Sci. Ser. Math. **61: 3** (1997), 75-90.
- [3] S. Bosch, U. Guntzer, R. Remmert. "Non-Archimedean analysis" (Berlin: Springer, 1984).
- [4] V.I. Bogachev, O.G. Smolyanov. "Analytic properties of infinite-dimensional distributions" // Usp. Mat. Nauk. **45: 3** (1990), 3-83.
- [5] N. Bourbaki. "Intégration". Livre VI. Fasc. XIII, XXI, XXIX, XXXV. Ch. 1-9 (Paris: Hermann, 1965, 1967, 1963, 1969) or (Moscow: Nauka, 1970 and 1977).
- [6] J.P.R. Christensen. "Topology and Borel structure". North-Holland Math. Studies N **10** (Amsterdam: Elsevier, 1974).
- [7] C. Constantinescu. "Spaces of measures" (Berlin: Springer, 1984).
- [8] Yu.L. Dalecky, S.V. Fomin. "Measures and differential equations in infinite-dimensional spaces" (Dordrecht: Kluwer Acad. Publ., 1991).
- [9] G.S. Djordjević, B. Dragovich. " p -Adic and adelic harmonic oscillator with a time-dependent frequency" // Theor. and Math. Phys., **124: 2** (2000), 1059-1067.
- [10] R. Engelking. "General topology" (Moscow: Mir, 1986).

- [11] H. Federer. "Geometric measure theory" (Berlin: Springer, 1969).
- [12] F.R. Gantmaher. "Theory of matrices" (Moscow: Nauka, 1988).
- [13] I.M. Gelfand, N.Ya. Vilenkin. "Some applications of harmonic analysis. Generalized functions". V. **4** (Moscow: Fiz.-Mat. Lit., 1961).
- [14] P.L. Henneken, A. Torta. "Theory of probability and some its applications" (Moscow: Nauka, 1974).
- [15] E. Hewitt, K.A. Ross. "Abstract harmonic analysis" (Sec. ed., Berlin: Springer, 1979).
- [16] C.J. Isham, J. Milnor in: "Relativity, groups and topology. II". P.1007-1290, Editors Witt B.S. De, Stora R.(Amsterdam: Elsevier, 1984).
- [17] Y. Jang. "Non-Archimedean quantum mechanics" // Tohoku Math. Publ. N **10**, 1998.
- [18] A.Yu. Khrennikov. "Mathematical methods of non-Archimedean physics"// Usp. Mat. Nauk. V. **45**: **4** (1990), 79-110.
- [19] A. Yu. Khrennikov, M. Endo. "Non-boundedness of p -adic Gaussian distributions"// Izvest. Akad. Nauk SSSR, Ser. Mat. **56** (1992), 1104-1115.
- [20] S.V. Ludkovsky "Measures on groups of diffeomorphisms of non-Archimedean Banach manifolds"// Uspekhi Mat. Nauk. **51**: **2** (1996), 169-170.
- [21] S.V. Ludkovsky. "Non-Archimedean polyhedral expansions of ultrauniform spaces"// Fundam. i Prikl. Mat., to appear.
- [22] S.V. Ludkovsky. "Measures on diffeomorphism groups of non-Archimedean manifolds, representations of groups and their applications"// Theor. and Math. Phys. **119**: **3** (1999), 381-396.
- [23] S.V. Ludkovsky. "Quasi-invariant measures on non-Archimedean semi-groups of loops"// Usp. Mat. Nauk. **53**: **3** (1998), 203-204.

- [24] S.V. Ludkovsky. "Irreducible unitary representations of non-Archimedean groups of diffeomorphisms" // Southeast Asian Bulletin of Mathematics. **22** (1998), 419-436.
- [25] S.V. Ludkovsky. "Properties of quasi-invariant measures on topological groups and associated algebras" // Annales Mathem. B. Pascal. **6: 1** (1999), 33-45.
- [26] S.V. Ludkovsky. "Quasi-invariant measures on non-Archimedean groups and semigroups of loops and paths, their representations" // Annales Mathematiques Blaise Pascal. **7: 2** (2000), 19-53, 55-80.
- [27] S.V. Ludkovsky. "Non-Archimedean free Banach spaces" // Fund. i Prikl. Mat. **1: 3** (1995), 979-987.
- [28] S.V. Ludkovsky. "Quasi-invariant measures on a group of diffeomorphisms of an infinite-dimensional real manifold and induced irreducible unitary representations" // Rendiconti dell'Istituto di Matem. dell'Università di Trieste. Nuova Serie. **31** (1999), 101-134.
- [29] A. Mađrecki. "Minlos' theorem in non-Archimedean locally convex spaces" // Comment. Math. (Warsaw). **30** (1991), 101-111.
- [30] A. Mađrecki. "Some negative results on existence of Sazonov topology in l -adic Frechet spaces" // Arch. Math. **56** (1991), 601-610.
- [31] A. Mađrecki. "On Sazonov type topology in p -adic Banach space" // Math. Zeit. **188** (1985), 225-236.
- [32] A.P. Monna, T.A. Springer. "Integration non-archimédienne" // Indag. Math. **25** (1963), 634-653.
- [33] L. Narici, E. Beckenstein. "Topological vector spaces" (New York: Marcel Dekker Inc., 1985).
- [34] A.C.M. van Rooij. "Non-Archimedean functional analysis" (New York: Marcel Dekker Inc., 1978).
- [35] W.H. Schikhof. "Ultrametric calculus" (Cambridge: Camb. Univ. Press, 1984).

- [36] W.H. Schikhov. "On p -adic compact operators". Report **8911** (Dept. Math. Cath. Univ., Nijmegen, The Netherlands, 1989).
- [37] W.H. Schikhof. "A Radon-Nikodym theorem for non-Archimedean integrals and absolutely continuous measures on groups" // Indag. Math. Ser. A. **33: 1** (1971), 78-85.
- [38] A.V. Skorohod. "Integration in Hilbert space" (Berlin: Springer, 1974).
- [39] O.G. Smolyanov, S.V. Fomin. "Measures on linear topological spaces" // Usp. Mat. Nauk. **31: 4** (1976), 3-56.
- [40] F. Topsøe. "Compactness and tightness in a space of measures with the topology of weak convergence" // Math. Scand. **34** (1974), 187-210.
- [41] F. Topsøe. "Some special results on convergent sequences of Radon measures" // Manuscripta Math. **19** (1976), 1-14.
- [42] N.N. Vahaniya, V.I. Tarieladze, S.A. Chobanyan. "Probability distributions in Banach spaces" (Moscow: Nauka, 1985).
- [43] V.S. Vladimirov, I.V. Volovich, E.I. Zelenov, " p -adic analysis and mathematical physics" (Moscow: Nauka, 1994).
- [44] A. Weil. "Basic number theory" (Berlin: Springer, 1973).

Address: Theoretical Department,
 Institute of General Physics,
 Russian Academy of Sciences,
 Str. Vavilov 38, Moscow, 119991 GSP-1, Russia